

DR. EMMA C LOCKLEY (Orcid ID : 0000-0003-2932-9945)

Article type : Special Issue Review and Syntheses

Effects of global warming on species with temperature-dependent sex determination: Bridging the gap between empirical research and management.

Emma C. Lockley¹ & Christophe Eizaguirre¹

¹School of Biological and Chemical Sciences, Queen Mary University London.

Abstract

Global warming could threaten over 400 species with temperature-dependent sex determination (TSD) worldwide, including all species of sea turtle. During embryonic development, rising temperatures might lead to the over-production of one sex, and, in turn, could bias populations' sex ratios to an extent that threatens their persistence. If climate change predictions are correct, and biased sex ratios reduce population viability, species with TSD may go rapidly extinct unless adaptive mechanisms, whether behavioural, physiological or molecular, exist to buffer these temperature-driven effects. Here, we summarize the discovery of the TSD phenomenon and its still elusive evolutionary significance. We then review the molecular pathways underpinning TSD in model species, along with the hormonal mechanisms that interact with temperatures to determine an individual's sex. To illustrate evolutionary mechanisms that can affect sex determination, we focus on sea turtle biology, discussing both the adaptive potential of this threatened TSD taxon, and the risks associated with conservation mis-management.

Keywords: Climate change, temperature-dependent sex determination, reptiles, sex steroid hormones, sea turtles

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/EVA.13226](https://doi.org/10.1111/EVA.13226)

This article is protected by copyright. All rights reserved

The current rate of species loss is often referred to as the sixth mass extinction event in geological history (Barnosky *et al.*, 2011). As climate change progresses and global temperatures continue to rise rapidly, understanding how species interact with their environment has become extremely important (Visser, 2008; Hoffmann *et al.*, 2011; Stocker *et al.*, 2013; Neukom *et al.*, 2019). This is particularly true for over 400 fish and reptile species that have temperature-dependent sex determination systems (TSD). For these species, incubation temperatures differentially trigger the pathways that lead to male and female gonad development (Charnov *et al.*, 1977; Deeming *et al.*, 1988). As global warming continues, the TSD mechanism could lead to heavily skewed offspring sex ratios towards one sex, which in turn threatens populations' persistence (Mitchell *et al.*, 2010; Laloë *et al.*, 2014; Eberhart-Phillips *et al.*, 2017). Whether TSD species will be able to withstand the rates of predicted temperature change and maintain viable sex ratios will depend on their adaptive potential (Eizaguirre *et al.*, 2014; Tomillo *et al.*, 2020). Adaptive potential is defined as "the ability of populations/species to respond to selection by means of phenotypic or molecular changes", and interacts with population structure and demography (Eizaguirre *et al.*, 2014; Rey *et al.*, 2020). Even though knowledge on the adaptive potential of TSD is essential for conservation management, it remains uncharacterised in most species, such as sea turtles (Tomillo *et al.*, 2020).

To understand pressures on sea turtles, it is essential to acknowledge that in addition to anthropogenic climate change, they face cumulative impacts from other human-induced stressors (McMahon and Hays, 2006; Hawkes *et al.*, 2007; Witt *et al.*, 2010), such as coastal development (Kaska *et al.*, 2013; Von Holle *et al.*, 2019), fisheries bycatch (Fossette *et al.*, 2014; Senko *et al.*, 2014) and illegal harvest of both eggs and adults (Tomillo *et al.*, 2008; Senko *et al.*, 2014). As a consequence, many populations are already depleted or in decline, and subject to extensive conservation management plans (Mortimer *et al.*, 2008; Hamann *et al.*, 2010; Wallace *et al.*, 2013). The actions to protect sea turtles from the effects of global warming are further limited by the inability to determine a neonate's sex non-lethally and the difficulty in justifying the sacrifice of individuals from endangered populations, both of which restrict TSD research in this taxon. As such, most interest in trying to quantify how sea turtles will adapt to climate change has focused on nesting behaviour such as phenological changes and site selection (Refsnider *et al.*, 2013; Mazaris *et al.*, 2013; Reneker and Kamal, 2016; Patrício *et al.*, 2017). Similarly, most approaches to mitigate the effects of rising temperatures have involved human manipulation of nest temperatures through, for example, relocation (either *in situ* or in hatcheries) and shading

(Mrosovsky, 2006; Tuttle *et al.*, 2010; DeGregorio *et al.*, 2011). Here, we review the current knowledge and highlight the importance of the adaptive potential of TSD mechanisms, by bridging empirical research gained from TSD model species with the more practical management of wild populations of sea turtles. We chose to highlight molecular and physiological responses, which we consider to be under-represented in sea turtle research in comparison to behavioural adjustments (Reneker *et al.*, 2016; Patrício *et al.*, 2017). Finally, we discuss how failing to consider the adaptive potential and underlying mechanisms of TSD in sea turtles could lead to inappropriate management decisions.

TSD patterns and environmental covariates

TSD species have no sex chromosomes. Instead, sex-determining genes are scattered across the genomes, and male- or female-determining pathways are triggered by temperature during a thermosensitive period of development (Charnov *et al.*, 1977; Bachtrog *et al.*, 2014; Shen *et al.*, 2014). This mode of sexual development was first reported in the common agama lizard, *Agama agama* in 1966 (Charnier, 1966). It has since been confirmed as the sex determining mechanism of several reptile lineages, including the tuatara, crocodilians and turtles (Janzen *et al.*, 1991; Cree *et al.*, 1995)

Different patterns of TSD exist (Figure 1). In Type Ia TSD, seen in most turtle species, males develop at cooler temperatures while females are produced under warmer conditions (e.g. the painted turtle, *Chrysemys picta*, Bull and Vogt, 1979). In Type Ib, this pattern is reversed, and males are produced at warm temperatures (e.g. the tuatara, *Sphenodon punctatus*, Cree *et al.*, 1995). Finally, species with Type II TSD, common to all crocodilians, produce males at intermediate temperatures and females at both hot and cold extremes (e.g. the American alligator, *Alligator mississippiensis*, Ferguson *et al.*, 1983; González *et al.*, 2019). Under constant incubation temperatures, the TSD thermal response curve is described by i) a pivotal temperature, at which an equal number of embryos within a clutch develop as males and females, and ii) the range of temperatures under which either male or female offspring may be produced, known as the transitional range of temperatures (Figure 1, Mrosovsky *et al.*, 1991; Girondot, 1999).

Genetic sex determination (GSD) and TSD are often considered to be mutually exclusive mechanisms, but instead should be regarded as two ends of a continuum, with environmental variation interacting with genetic mechanisms to different extents across species (Quinn *et al.*, 2007, 2011; Pen *et al.*, 2010; Bachtrog *et al.*, 2014; Holleley *et al.*, 2015, 2016). For instance, the

94 montane lizard, *Bassiana duperreyi*, has heteromorphic sex chromosomes, and eggs incubated
95 at warm temperatures, characteristic of lowland environments, produce equal numbers of male
96 and female offspring. Yet, when eggs are incubated at cool temperatures representative of high
97 altitude summers, the ratio is skewed towards 70% male hatchlings, suggesting that temperature
98 can partly override genetic triggers in this species (Shine *et al.*, 2002). In mammals, which have
99 GSD, maternal condition can alter sex ratios (Trivers *et al.*, 1973; Sheldon *et al.*, 2004; Grant,
100 2007), moderated by the environmental conditions during the breeding and gestation periods
101 (Edwards *et al.*, 2019).

102 Temperature is the primary determinant of gonad differentiation in TSD species, and the thermal
103 environment of nests varies with substrate albedo (Hays *et al.*, 2001), shading (Refsnider *et al.*,
104 2013) and nest depth (Telemeco *et al.*, 2009; Tomillo *et al.*, 2017), which can all influence egg
105 development. These variables differ depending on individual nest site selection and seasonality,
106 and thus introduce variation into offspring sex ratios within and among populations (Reneker &
107 Kamal, 2016). It is also known that precipitation and humidity interact with thermal conditions, and
108 hence influence TSD. For instance, the relationship between rainfall and sex ratios has generally
109 been attributed to the cooling effect of rain on the temperature of nesting substrate thanks to
110 evaporative cooling (Godfrey *et al.*, 1996; Matsuzawa *et al.*, 2002; Houghton *et al.*, 2007; Lolavar
111 *et al.*, 2015). An argument has recently also been made to suggest that humidity itself has an
112 effect on sex ratios beyond that of temperature alone (Lolavar *et al.*, 2017, 2020). While these
113 studies have not yet robustly demonstrated the effect of humidity on TSD, they do illustrate
114 several points. Firstly, temperature does not act in isolation in natural environments. Secondly,
115 we still do not fully understand all the factors that influence of sex determination in TSD species.
116 As such, we should consider the potential for elements of the thermal response curve (e.g. the
117 pivotal temperature and transitional range of temperatures) to vary from traditional values - either
118 from plastic responses or as a result of adaptive evolution (Santidrián Tomillo *et al.*, 2020).

119 ***The Evolutionary Significance of TSD***

120 A comprehensive theory explaining the selective advantage of TSD still seems to evade
121 researchers (Sarre *et al.*, 2004; Pen *et al.*, 2010; Quinn *et al.*, 2011). While the random
122 segregation of sex chromosomes in species with GSD reflects parents' equal chromosomal
123 investment in male and female offspring, conforming to frequency-dependent selection, no such
124 theory exists to easily explain the evolutionary significance of TSD (Fisher, 1930). Instead, three
125 relatively robust hypotheses for the evolution of TSD have been suggested, with varying levels of
126 support; i) the Charnov-Bull model of differential fitness, ii) the "Mighty Males" hypothesis, and iii)

127 the phylogenetic inertia hypothesis (Charnov *et al.*, 1977; Girondot *et al.*, 1999; Shine, 1999;
128 Janzen *et al.*, 2006; Rollinson, 2019).

129 The Charnov-Bull model of differential fitness suggests that sex-specific advantages are
130 associated with particular thermal environments, and that the TSD mechanism ensures the
131 production of sexes at their optimal temperatures (Charnov *et al.*, 1977). The theory requires a
132 heterogeneous environment, in time or space, where males and females benefit from different
133 conditions (Charnov *et al.*, 1977). Sex-specific benefits from particular thermal environments
134 emerge through different mechanisms. For instance incubation temperature may i) affect fitness
135 proxies, such as growth rate or size, ii) correlate with future conditions that offspring will
136 experience, or iii) affect developmental rates and timing of emergence (Shine, 1999; Janzen *et al.*, 2006). Recently it was suggested that sex-specific fitness may stem from bimodal age at
137 maturity distributions, as TSD species show greater dimorphism in this trait than GSD species
138 (Bókony *et al.*, 2019).

140 In TSD species the effects of temperature and sex overlap, and therefore it is challenging to
141 disentangle their relative contributions to an individual's fitness (Rhen *et al.*, 2004). This problem
142 has generally been overcome with the use of exogenous hormone manipulation experiments,
143 whereby a given sex is artificially forced to develop at temperatures that would not otherwise
144 enable its production. One of the best examples of such an experiment focused on the jacky
145 dragon, *Amphibolurus muricatus* (Warner *et al.*, 2008). Eggs from this agamid lizard were
146 incubated at a range of temperatures, and half of them were treated with an aromatase inhibitor
147 that forced embryos to develop as male, regardless of their thermal environment. This
148 manipulation decoupled the effects of sex and temperature on fitness and revealed that lifetime
149 reproductive success was greater for males that were incubated at natural male-producing
150 temperatures (Warner *et al.*, 2008). Importantly, hormone treatment had no effect on the
151 morphology or survivorship of juvenile jacky dragons, indicating no effect of the treatment itself.
152 Since this seminal study, further experiments have produced similar results across other agamid
153 species (Steele *et al.*, 2020). While this reproducibility gives weight to the Charnov-Bull
154 hypothesis, whether these results can be extrapolated to all TSD species, and particularly long-
155 lived ones, remains unclear (Steele *et al.*, 2020). This is because there are conflicting results
156 among experiments testing the Charnov-Bull model (Janzen *et al.*, 2006). For instance, studies
157 using the diamondback terrapin *Malaclemys terrapin* failed to find support for the Charnov-Bull
158 model (Morjan *et al.*, 2003). As such, caution is needed when presenting the Charnov-Bull model
159 as a universal explanation for TSD in reptiles (Janzen *et al.*, 2006).

160 Recently, Rollinson (2019) proposed an alternative theory for the evolution of TSD, referred to as
161 the “Mighty Males” hypothesis, based on the maternal condition hypothesis described by Trivers
162 and Willard (1973). The original hypothesis posits that females’ lifetime reproductive success is
163 mostly constrained by the number of gametes they produce and, as such, even lower quality
164 female offspring produced under sub-optimal conditions should not suffer a reduction in lifetime
165 reproductive fitness. On the other hand, males’ reproductive success is limited by their ability to
166 compete for mating opportunities, and therefore male offspring should be produced under the
167 environmental conditions that maximise their overall fitness (Trivers *et al.*, 1973). Given these
168 constraints are related to sexual reproduction, independently of TSD, and given that temperature
169 affects a wide range of traits such as size, developmental rate, or embryonic mortality, Rollinson
170 (2019) proposes that males should be produced at the temperatures that maximise the fitness
171 potential of these traits, while females should be produced at the sub-optimal extremes. As
172 mortality increases with high temperatures, the Mighty Males hypothesis can be easily applied to
173 type II (female-male-female) and type Ia (male-female) TSD species (Santidrián Tomillo *et al.*,
174 2020). On the other hand, it may not apply to type Ib (female-male) TSD species, unless
175 evolutionary mechanisms exist that can significantly push back the upper temperature limits for
176 mortality. Type Ib is rare in nature, existing in the tuatara and shorter-lived squamata groups, and
177 may require an alternative explanation.

178 Finally, the difficulty in determining the evolutionary significance of TSD may stem from
179 phylogenetic inertia and the possibility its adaptive significance may no longer be detectable 300
180 million years after it evolved (Janzen *et al.*, 2004, 2006). After phylogenetically reconstructing the
181 evolution of sex determination in over 400 species of squamata, many examples of transitions
182 from TSD to GSD were identified, but there were no cases where this direction was reversed
183 (Pokorná *et al.*, 2009). It was therefore suggested that sex chromosomes may evolve when sex-
184 specific genes are coupled with genes that provide a selective advantage to that sex. As the
185 association between these genes strengthens and rates of recombination decrease, the
186 conditions are met for sex chromosomes to evolve (Bachtrog *et al.*, 2014; Muralidhar *et al.*,
187 2018).

188 ***Heritability, Development and Molecular Pathways of Sex Determination.***

189 Despite very different sex determination mechanisms across vertebrate taxa, similar genetic
190 pathways underpin the molecular foundations of gonad differentiation. This is consistent with
191 conserved patterns of gonad development across vertebrates, with gonads being initiated as a
192 bipotential genital ridge in both GSD and TSD species. From a developmental perspective, the

193 genital ridge is formed from the coelomic epithelium, underlying mesenchymal cells and germ
194 cells, which migrate into the ridge from the embryonic yolk sac (Morrish *et al.*, 2002). As the
195 somatic and germ cells proliferate, the genital ridge expands until the cell fate commits to
196 developing as testes or ovaries. The differentiation of testes necessitates the development of the
197 primary sex cords into testis cords, whereas the formation of ovaries requires the thickening of
198 the coelomic epithelium while the primary sex cords disappear (Morrish *et al.*, 2002). Importantly,
199 depending on the species and the TSD pattern, the molecular mechanisms of the sex-
200 determining period can either end before signs of gonad differentiation appear, or may overlap
201 with the early stages of gonad differentiation (Morrish *et al.*, 2002).

202 Overall, there is no doubt about the role of temperature in sex determination of TSD species, yet
203 individual responses, reflected in traits such as pivotal temperature, are underpinned by a genetic
204 contribution with some levels of heritability, estimated in the painted turtle, *Chrysemys picta*, to be
205 as high as $h^2=0.35$ (McGaugh *et al.*, 2011). It is therefore not surprising that studies have focused
206 on elucidating the molecular mechanisms of sex determination (Martínez-Juárez *et al.*, 2019).
207 Conventionally, genes are classified based on their GSD sex-specific pathways and probably all
208 known mammalian sex determining genes have now been tested for their role in TSD (Rhen and
209 Schroeder 2010). A very thorough gene-based overview can be found in Martínez-Juárez &
210 Moreno-Mendoza (2019). In brief, the expression of genes such as *Foxl2* (Forkhead box protein
211 L2) and *Rspo1* (encoding the protein R-spondin-1) becomes greater in the female determining
212 pathway, but these are also detectable in the male cascade during early development (Elf 2003).
213 Furthermore, *Dax1* (dosage-sensitive sex reversal, adrenal hypoplasia critical region), *Sf1*
214 (steroidogenic factor 1) and *Wnt4* (Wingless-Type MMTV Integration Site Family, Member 4), all
215 involved in female development and the repression of male traits, also appear to contribute to the
216 sex determination of TSD species (Shoemaker and Crews 2009). Conversely, in the male
217 pathway, genes necessary for testicular differentiation include *Sox9* (SRY-Box Transcription
218 Factor 9), the anti-Müllerian hormone (*AMH*) and *Dmrt1* (Doublesex And Mab-3 Related
219 Transcription Factor 1). The expression of *Sox9*, a gene that belongs to the same HMG-box
220 transcription factor as *Sry* (Sex determining region Y), is detectable in early gonad development
221 and becomes restricted to the developing testis at the end of the thermosensitive period
222 (reviewed in Rhen and Schroeder 2010). Noteworthy, the expression of *Sry* remains elusive in
223 reptiles. *Dmrt1* is particularly interesting and shows temperature-dependent sex-specific
224 expression that precedes gonadal sex differentiation. *Dmrt1* expression changes with shifts in
225 temperature in a sex-specific manner, and also responds to the presence of aromatase inhibitors
226 – regulation of this gene has been shown to be necessary in order to initiate male development in

227 *Trachemys scripta* (Ge *et al.*, 2017).

228 While the identification of these genes follows a candidate gene approach, in more recent years
229 studies of TSD have focused on gene discovery. Along this vein, several new candidates have
230 emerged. A single nucleotide polymorphism in *Cirbp* (Cold Inducible RNA Binding Protein) was
231 associated with transcript levels in the embryonic gonads of the snapping turtle (*Chelydra*
232 *serpentina*) during specification of gonad fate and hatchling sex (Schroeder *et al.*, 2016; Rhen *et*
233 *al.*, 2017). The A allele was induced in embryos exposed to a female-producing temperature,
234 while expression of the C allele did not differ between female- and male-producing temperatures.
235 As such AA homozygotes were more likely to develop ovaries than the CC homozygotes which
236 all developed as males, with the AC heterozygotes standing at an intermediate frequency. From
237 an ecological perspective, it is worth noting that changes in allele frequencies in *Cirbp* were
238 detected at small and large geographical scales, suggesting local adaptation. Such patterns of
239 local adaptation would be expected to result in higher pivotal temperatures in warmer regions,
240 possibly mediated by resistance thresholds to the development of the high temperature sex,
241 linked to genetic underpinnings. On the other hand, in the central bearded dragon (*Pogona*
242 *vitticeps*), a species in which chromosomal sex determination is overridden at high temperatures,
243 sex-reversed females are produced when an intron is retained in the mature transcripts from
244 each of two Jumonji family genes, *Jarid2* (Jumonji And AT-Rich Interaction Domain Containing 2)
245 and *Jmjd3* (Jumonji Domain Containing 3, Histone Lysine Demethylase). This intron retention
246 was observed only in females that have been sex-reversed by temperature, not in classic
247 chromosomal females or males (Deveson *et al.*, 2017). Similarly to *Cirbp* in *C. serpentina*, if the
248 central bearded dragon has evolved mechanisms to regulate the over production of a given sex,
249 we should be able to observe spatial and temporal variation of this intron loss in relation to
250 temperature clines.

251 De-novo genome sequencing has also introduced elements of gene and mechanism discovery.
252 This has been the case with the assembly of American alligator genomes, which, combined with
253 RNA sequencing and models of CTCF-mediated chromatin looping, identified genomic regions
254 that were significantly enriched for genes with female-biased expression in developing gonads
255 after the thermo-sensitive period (Rice *et al.*, 2017). This approach demonstrated that estrogen
256 signalling is a major driver of female-biased gene expression and holds promises for comparative
257 genomics of TSD species.

258 Finally, a common pattern that has emerged from genomic studies is that the discovery of new
259 TSD associated genes does not appear to be independent of epigenetic mechanisms. For

instance, the DNA methylation dynamics of the *Dmrt1* promoter region are tightly correlated with temperature and could mediate the impact of temperature on sex determination (Ge *et al.*, 2018). In *T. scripta*, Ge et al (2018) demonstrated how the epigenetic regulator *Kdm6b* demethylates the histone H3 lysine 27 (*H3K27*) at the *Dmrt1* promoter region, in a process that results in male sex determination. It appeared though that *Kdm6b* is not in itself responsive to temperature, and as such the fundamental thermal trigger of this pathway remains unknown. Similarly, *Jarid2* is a component of the master chromatin modifier Polycomb Repressive Complex 2, and the mammalian sex-determining factor *Sry* is directly regulated by an independent but closely related Jumonji family member (Devison et al 2017). The authors proposed that the alteration of *Jarid2/Jmjd3* function by intron retention alters the epigenetic landscape to override chromosomal sex-determining cues, triggering sex reversal at extreme temperatures (Devison et al 2017).

It is difficult to predict the nature of selection exerted by global warming, as directional selection would be expected under a constant increase in temperature, but a predicted increased frequency of extreme events should result in fluctuating selection. When selection is constrained, then phenotypic plasticity may allow species to respond to their environment (Chevin *et al.*, 2010). Indeed, while epigenetics, as mechanisms of phenotypic plasticity, may appear distant to practical conservation, their relevance has recently been framed in the light of management decisions (Rey *et al.*, 2020). Altogether, understanding the mechanisms underlying phenotypic plasticity is a valuable direction of research in conservation management. In particular, here we suggest that increased consideration of endocrinology and sex steroid hormones would benefit the conservation of TSD species.

Plasticity and sex steroid hormones in TSD species

Physiological plasticity includes a series of mechanisms by which organisms can match their phenotypes to their environmental conditions, for instance via the endocrine system and hormone regulation (Gienapp *et al.*, 2008; Chevin *et al.*, 2010; Merilä *et al.*, 2014). In the specific context of phenotype-environment matching, evolutionary theory suggests that the populations demonstrating the greatest levels of plasticity in beneficial traits might have the highest potential of persisting in the face of climate change (Meyers *et al.*, 2002). This is because if the cost of plasticity is low, it will reduce the range of conditions under which extinction is inevitable (Chevin *et al.*, 2010; Sanford *et al.*, 2011). Plasticity could thus maintain populations until adaptive evolution emerges and improves the phenotype-environment match (Lande, 2009; Reusch, 2014).

292 Sex steroid hormones, such as estrogens and their precursors, androgens, interact with the
293 molecular pathways that control gonad differentiation, and are fundamental for regulating gonad
294 development across a wide variety of taxa (spanning amphibian (Ko *et al.*, 2008), bird
295 (Nakabayashi *et al.*, 1998), fish (Wang *et al.*, 2007), mammals (Uhlenhaut *et al.*, 2009) and
296 reptiles (Barske *et al.*, 2010)). Estrogens repress *Sox9* expression, preventing the differentiation
297 of male gonad-specific Sertoli cells in both GSD (mouse, Uhlenhaut *et al.* 2009) and TSD species
298 (*T. scripta*, Barske and Capel, 2010). Conversely, the development of the müllerian ducts, which
299 provide the structure for female gonads, correlates with changes in estrogens in mammals, birds
300 and reptiles (Dodd *et al.*, 2008). In TSD species, the estrogen 17 β -estradiol replicates the effect
301 of temperature in instigating demethylation of histone H3 lysine 27 (*H3K27*) at the *Dmrt1*
302 promoter region, resulting in female gonad development (Ge *et al.*, 2018). Moderating embryonic
303 exposure to sex steroid hormones is thus likely to be an effective mechanism for conferring
304 plasticity to the TSD thermal response curve (Bowden *et al.*, 2000; Carter *et al.*, 2017).

305 The influence of androgens and estrogens on embryonic sex independently of temperature in
306 TSD species has been extensively reviewed (Elf, 2003; Bowden and Paitz, 2018). To date,
307 research exploring the relationship between steroid hormones and TSD pathways has largely
308 focused on estradiol, testosterone and the enzyme aromatase (Box 1). Studies began with *in vitro*
309 manipulation experiments which showed that exogenous application of estradiol produced female
310 offspring at male-producing temperatures (Crews *et al.*, 1989, 1991; Wibbels *et al.*, 1991). At the
311 same time, the application of aromatase inhibitors such as fadrozole, which prevent the synthesis
312 of estradiol, have repeatedly resulted in male offspring (Wibbels *et al.*, 1994; Rhen *et al.*, 1995;
313 Warner *et al.*, 2017). Patterns of gonad development in response to exogenous hormone
314 application have not, however, always been predictable and negative results have been reported,
315 along with incidences where exogenous application of estradiol unexpectedly produced male
316 offspring (Janes *et al.*, 2007; Warner *et al.*, 2014).

317 The apparent inconsistencies that are reported both between *in vitro* experiments and across
318 species can be partially explained by interactions between exogenous hormones and
319 temperature. Estradiol and temperature produced more female embryos than would be expected
320 by each treatment alone near the pivotal temperature of the red-eared slider turtle *T. scripta*
321 (Wibbels *et al.*, 1991). Under extreme natural incubation temperatures (>2 °C above the 75 year
322 nesting site average), exogenous application of estradiol and fadrozole had no effect on painted
323 turtle sex ratios (Warner *et al.*, 2017). Yet, under average seasonal conditions exogenous
324 estradiol produced more female offspring, and fadrozole more male offspring, than controls

325 (Warner *et al.*, 2017). These results suggest that temperature and hormones may have a dose-
326 dependent and interactive effect on sex determination pathways (Wibbels *et al.*, 1991).

327 In oviparous species, a principal conduit of hormone transfer from mother to offspring is the egg
328 yolk (Schwabl, 1993; Radder, 2007). It provides material with which to prime the reactions
329 associated with the molecular cascades that trigger gonad differentiation. In European sea bass
330 *Dicentrarchus labrax*, exposure to male producing temperatures results in methylation of the
331 *Cyp19a1* gene and lower aromatase expression (Navarro-Martín *et al.*, 2011). Similar methylation
332 patterns are also seen in red-eared slider turtles *T. scripta* (Matsumoto *et al.*, 2016). High levels
333 of aromatase expression will increase the biosynthesis of estradiol from testosterone, if this
334 substrate is available and the reaction is not inhibited (Boon *et al.*, 2012). Interestingly, the
335 estradiol:testosterone ratio within egg yolks at oviposition (i.e. before synthesis has occurred)
336 varies across reptile species, having been recorded as both above and below 1, but rarely
337 spanning an equal ratio (Radder, 2007), suggesting species-specific regulation. Depending on
338 species, endogenous estradiol in yolks could act on female-producing pathways directly, or
339 indirectly through the aromatase synthesis of endogenous testosterone, to produce female
340 hatchlings (Figure 2). However, in loggerhead sea turtles, the ratio of estradiol:testosterone in
341 egg yolks can favour either hormone, or be equal (Lockley *et al.* 2020). In a natural experiment
342 where temperatures were controlled among nests, equal, low concentration,
343 estradiol:testosterone ratios produced male offspring, which the authors theorise is due to
344 product-feedback inhibition of aromatase (Figure 2, Lockley *et al.* 2020). Such reactions will be
345 occurring within the yolk of eggs used for exogenous application studies. The interference with
346 natural estradiol:testosterone ratios that could occur in these experiments might thus explain
347 those occasions where exogenous estradiol application has produced male offspring (Janes *et*
348 *al.*, 2007; Warner *et al.*, 2014).

349
350 A fundamental constraint of our knowledge generated by exogenous application studies is
351 therefore that the biological relevance of results within natural systems is extremely limited
352 (Bowden *et al.*, 2018). Field studies are invaluable for elucidating how this mechanism interacts
353 with environment. When clutches of painted turtles were incubated at constant temperatures of
354 28 °C, the endogenous estradiol:testosterone ratio within egg yolks correlated with a seasonal
355 shift in sex ratio from 72% male to 76% female (Bowden *et al.*, 2002). This relationship resulted in
356 thermal reaction curves systematically changing across a nesting season, in a manner that
357 allowed sex-specific phenotypic matching, enhancing the production of female offspring under
358 warm conditions (Carter *et al.*, 2017). Such a mechanism fits directly with the Charnov-Bull

359 differential fitness hypothesis, allowing nesting females to maximise female offspring production
360 under temperatures that could be most beneficial to them. In addition, it is consistent with the
361 “Mighty Males” hypothesis, favouring the production of female offspring under warm conditions
362 that may increase mortality rates. This mechanism, however, could accelerate the rate of female
363 production as temperatures rise and may therefore represent an example of evolutionary suicide
364 (Bowden *et al.*, 2000).

365 ***Temperature-dependent Sex Determination and Sea Turtles***

366 We have considered the many molecular pathways that contribute to the formation of gonads in
367 TSD species, and outlined how sex steroid hormones can interact with temperature (Figure 3).
368 We now focus on applying this gained knowledge to sea turtles and their management, in the
369 light of global warming. Sea turtles are Type Ia TSD species (Yntema *et al.*, 1982) and thus
370 thermal projections across the coming century suggest that offspring sex ratios will become
371 increasingly feminised (Yntema & Mrosovsky 1982; Hawkes *et al.* 2007; Hawkes *et al.* 2009; Witt
372 *et al.* 2010; Laloë *et al.* 2014; Tanner *et al.* 2019). These projections, however, do not generally
373 account for the adaptive potential of populations (Santidrián Tomillo *et al.*, 2020).

374 Sea turtles demonstrate a high degree of natal philopatry to their nesting sites, often with high
375 female fidelity and male-mediated gene flow among mating sites (Meylan *et al.*, 1990; Bowen *et al.*
376 *et al.*, 2007; Lee *et al.*, 2007; Stiebens *et al.*, 2013; Levasseur *et al.*, 2020). This strong female
377 philopatry creates genetic structure within a rookery that is prone to the evolution of local
378 adaptation (Stiebens *et al.*, 2013; Baltazar-Soares *et al.*, 2020). Female turtles return to nest at
379 intervals ranging from 1 to 4 years, while males return more frequently, and can even remain
380 resident to the mating grounds (Hays *et al.*, 2010; Schofield *et al.*, 2010; Arendt *et al.*, 2012).
381 Different re-migration intervals between males and females result in different operational sex
382 ratios (the ratio of sexually active males to females at the nesting site a given time) than that of
383 adult sex ratios, which are likely to be less female biased than offspring sex ratios (Hays *et al.*,
384 2014). While the operational sex ratio and even mating strategies (e.g. Lee *et al.*, 2004) may
385 alleviate offspring sex ratio bias, a reduction in the total number of males can have implications
386 for populations: it can lead to reduced genetic diversity (Frankham, 2005), increase potential for
387 inbreeding and genetic drift (Hedrick *et al.*, 2000), and ultimately reduce a population’s fitness
388 and its adaptive potential (Reed *et al.*, 2003).

389 ***TSD in Sea Turtles: The barriers***

390 Little is known about TSD in sea turtles as, historically, sexing individuals has required sacrifice
391 and histological examination of gonads (e.g. Wyneken *et al.*, 2007; Hamann *et al.*, 2010; Fuentes
392 *et al.*, 2017). As most sea turtle populations are listed on the IUCN red list, the sacrifice of
393 hatchlings is often restricted by local authorities. Over the years, several protocols have been
394 developed to overcome this challenge, such as quantifying the estradiol:testosterone ratio in
395 hatchling plasma (green sea turtle: 96.7% overall accuracy, $n = 30$, Xia *et al.*, 2011, loggerhead
396 sea turtle: 94% accuracy for males and 100% accuracy for females, $n = 28$, Gross *et al.*, 1995).
397 Along the same approach, the expression of Anti-Müllerian Hormone (*AMH*) seems to also
398 effectively identify the sex of loggerhead sea turtle hatchlings, being expressed only in males ($n =$
399 59, Tezak *et al.*, 2020). These protocols will now need to be up-scaled to reach a high
400 throughput, to lead a new era of research in the study of TSD and management of sea turtles.

401 Until large scale sexing techniques are possible, our overall knowledge of TSD thermal response
402 curves in sea turtles is limited to laboratory incubation studies with small sample sizes (Rimblot *et al.*,
403 1985; Mrosovsky, 1988; Mrosovsky *et al.*, 1992; Godfrey *et al.*, 1999). These studies mostly
404 identified the pivotal temperature of sea turtle populations to lie around 29 °C (but see McCoy *et al.*
405 1983; Wibbels, *et al.*, 1998). In more recent years, sophisticated modelling approaches have
406 been used to refine the characteristics of thermal response curves (Abreu-Grobois *et al.*, 2020),
407 and have also quantified the variation in this trait among populations (Bentley *et al.*, 2020).
408 However, prior to these advances, 29 °C was often used as an approximate pivotal temperature
409 to estimate sex ratios of sea turtle populations in cases where sacrificing hatchlings was not
410 possible, using mean nest temperature during the middle third of incubation as a proxy (see
411 Table 1 for illustrative cases). Alternatively, studies have used the thermal response curves
412 described specifically for the population of interest or a nearby neighbour, if these values have
413 been reported (Table 1).

414
415 The approach of using a pivotal temperature proxy during the middle third of incubation is now
416 widely regarded to be too oversimplified, as the thermosensitive period is dependent on
417 development rates, which are not linearly related to temperature, particularly under variable
418 thermal environments (Georges *et al.*, 2005; Massey *et al.*, 2019). Importantly, it also does not
419 account for individual variation in, or the adaptive potential of the pivotal temperature. Other
420 measures that correlate with sex ratio, such as incubation duration, have also been used, but
421 estimates tend to vary dependent on the proxy used (Fuentes *et al.*, 2017). For instance, Fuentes
422 *et al.* (2017) demonstrated that estimating the sex ratio of natural nests using the constant
423 temperature equivalent (which converts natural fluctuating incubation temperatures into a

constant value to be compared to temperatures produced under lab conditions (Georges *et al.*, 2005)) during the middle third of incubation, and assuming a pivotal temperature of 29.12 °C, predicted an average sex ratio of 5.99% (95% CI: <0.1% - 37%) male. This estimate increased to 9.97% (95% CI: <0.1% - 86.20%) male when using incubation duration as a proxy for the same nests. While the use of proxies has contributed to our general understanding of sex ratio distributions, their reliance on pre-existing thermal response curves make them unable to detect real-time deviation from such curves that might originate from different elements of the adaptive potential of sea turtles – conceivably with major consequences of past and future management decisions.

433

434 ***TSD in Sea Turtles: Adaptive Potential***

To date, attention on the impact of global warming on the sex ratios of sea turtles has largely focused on how plastic behaviours can avoid extreme sex ratio biases. Such behaviours include nest site selection (Reneker & Kamal, 2016; Patrício *et al.*, 2017), nest depth (Refsnider *et al.*, 2013), and phenological shifts (Mazaris *et al.*, 2013). For instance, with rising temperatures there is evidence that nesting seasons start earlier and can be more protracted for loggerhead turtles nesting in North Carolina and Greece (Weishampel *et al.*, 2004; Hawkes *et al.*, 2007; Mazaris *et al.*, 2013; Patel *et al.*, 2016). From a global perspective, there is a significant negative relationship between the dates of first nesting for populations of loggerhead sea turtles across their nesting distribution, and the sea surface temperature at the beginning of the nesting season (Mazaris *et al.*, 2013). Despite evidence of phenological variation, the sole extent of such shifts is unlikely to be sufficient to keep up with the rate of contemporary climate change (Telemeco *et al.*, 2013; Monsinjon *et al.*, 2019).

447

Consequently, mechanisms other than behavioural adjustments may prove vitally important for sea turtles to respond to climate change. Local adaptation, which, in the case of TSD, would involve populations adjusting thermal response curves to match the changing thermal environment of their nesting sites, could be a fundamental part of this process. Patterns of local adaptation have been shown to evolve from the philopatric nature of sea turtles, as previously demonstrated for immune genes (Stiebens *et al.* 2013) and even feeding strategies (Cameron *et al.* 2019). For instance, green sea turtle hatchlings from dark sand (high temperature) beaches on Ascension Island grow faster and have higher levels of hatching success than those from nearby white sand (cool) beaches when exposed to hot artificial incubation environments (Weber *et al.*, 2012). Interestingly, no fine-scale adaptation of the pivotal temperature was found between turtles nesting on these two beaches in a later study, showing the context-dependence of the

459 results (Tilley *et al.*, 2019). On the other hand, flatback turtle *Natator depressus* eggs at tropical
460 latitudes in Australia have high levels of tolerance to prolonged warming exposure up to 35 °C
461 during incubation, despite these temperatures often being lethal (Maulany *et al.*, 2012; Howard *et*
462 *al.*, 2015). The pivotal temperatures of three genetically distinct flatback turtle populations also
463 vary by 1.5 °C, being highest in the population nesting under the warmest conditions (Bentley *et*
464 *al.*, 2020). These records of adaptation to local thermal environments provide evidence that the
465 temperature sensitive triggers of the molecular pathways controlling sex determination can evolve
466 (e.g. Ewert *et al.*, 2005).

467 There is very little information on whether maternally derived sex steroid hormones can contribute
468 to local adaptation, or whether they can introduce plasticity to the sea turtle TSD
469 mechanism. This lack of attention to the evolvability of the endocrine system is unwarranted
470 given the evidence from other TSD species (Bowden *et al.*, 2000; Ewert *et al.*, 2005).
471 Experimental estradiol treatment of olive ridley turtle eggs at male producing temperatures can
472 feminize gonads (Merchant-Larios *et al.*, 1997), disrupt testis differentiation (Díaz-Hernández *et*
473 *al.*, 2015), reduce cell proliferation (Díaz-Hernández *et al.*, 2017), delay Sox9 inhibition, and delay
474 the upregulation of *FoxL2* and aromatase (Díaz-Hernández *et al.*, 2015). Furthermore, emerging
475 research from a natural, *in situ* experiment that controlled for temperature found that the ratio of
476 maternally derived estradiol:testosterone in the yolk of loggerhead sea turtle eggs correlated with
477 the sex ratio of nests independently of temperature (Lockley *et al.*, 2020). Research that aims to
478 evaluate the elements of the TSD mechanism that can evolve and locally adapt will be
479 fundamental for revealing the adaptive potential of the TSD mechanism in sea turtles.

480 ***Implications for Conservation Management***

481 Global efforts dedicated to the conservation of sea turtles are extensive, with hundreds of
482 grassroot, national and international projects working to stop the decline of populations and
483 promote their persistence. Such work frequently focuses on the protection of nesting beaches
484 (Hamann *et al.*, 2010; Fuentes *et al.*, 2012). Along with the benefits of accessibility, logistical
485 advantages and low cost options, protection at this point in the life cycle can reduce threats from,
486 for example, poaching of eggs and adult turtles (Tomillo *et al.*, 2008; Senko *et al.*, 2014), tidal
487 inundation (Varela *et al.*, 2019) and coastal development (Kaska *et al.*, 2013; Von Holle *et al.*,
488 2019).

489 A common conservation approach to mitigate the effects of human-induced stressors such as
490 coastal development and poaching is to relocate egg clutches into *in situ* hatcheries, where nests

491 are protected and monitored (Mrosovsky, 2006, 2008; Pike, 2008; Pfaller *et al.*, 2009; Tuttle *et*
492 *al.*, 2010). However, these relocations should not be performed lightly, as shading, substrate
493 properties and depth can be different from the nesting beach, and can all substantially alter the
494 local thermal regimes (Morreale *et al.*, 1982; Tuttle *et al.*, 2010; DeGregorio *et al.*, 2011). The
495 differences between temperatures in hatcheries and *in situ* sea turtle nests vary between
496 locations. For instance, no difference in temperature was recorded during the thermosensitive
497 period between relocated and *in situ* loggerhead sea turtle nests in North Carolina, but over the
498 entire incubation period nests in the hatchery were exposed to higher overall temperatures, and
499 hatchlings from these nests emerged sooner (DeGregorio *et al.* 2011). However, this was not
500 observed in Georgia, where there were no significant differences in temperatures, size or
501 incubation duration between relocated and *in situ* nests, but there was reduced survival in the
502 nests that had been relocated (Tuttle *et al.*, 2010). If nest relocation is undertaken, then
503 continuous evaluation of incubation temperatures, durations and the effects on hatchling
504 development and fitness should always be conducted.

505 Crucially, if done incorrectly, relocation may interfere with natural selection for traits that will help
506 sea turtles adjust to new temperature regimes, or reduce the effectiveness of natural buffering
507 mechanisms, as climate change progresses (Mrosovsky, 2006; Pfaller *et al.*, 2009). This effect
508 may be particularly strong for these highly philopatric species, as they are the ones predicted to
509 have evolved the strongest signature of local adaptation (Stiebens *et al.*, 2013; Baltazar-Soares
510 *et al.*, 2020). Specifically, human manipulation of nest temperatures might dampen the effects of
511 natural selection on developing embryos, by reducing selection on the genomic mechanisms that
512 confer thermal tolerance and a higher pivotal temperature. Alternatively, if plastic phenotype-
513 environment matching were occurring, either in the form of maternal hormone transfer or nesting
514 behaviour, intentionally modifying incubation conditions would cause a human-induced thermal
515 mismatch between the clutch and its optimal conditions.

516 It is thus important that if nest relocation is necessary to protect from stressors such as
517 inundation or predation, this is done in a manner that matches the thermal environment that
518 embryos would have experienced had eggs remained *in situ*. This includes taking all possible
519 care to relocate nests to similar substrates in the vicinity of the nesting beach, in a nest cavity of
520 the same depth and shape as originally dug by the female, with similar hydric properties. By
521 doing this, conservationists will ensure that any behaviour is accounted for, while ensuring that
522 unseen physiological plasticity such as hormone transfer will act in the direction of natural
523 selection. In addition, in light of our current lack of knowledge on accurate sex ratios and adaptive

524 potential, relocating nests with the direct intention of manipulating temperature regimes (in an
525 attempt to mitigate against global warming itself) risks preventing populations from adapting to
526 conditions naturally. This leaves conservationists in a position where leaving egg clutches
527 exposed to increasing temperatures is not viable, but altering incubation conditions by reducing
528 temperatures may also have long-term negative consequences, whereby local adaptation is
529 prevented or reduced. For scientists interested in TSD, focusing on developing non-lethal sexing
530 methods that can be applied at a large scale would be extremely useful to guide such future
531 conservation decisions.

532 **Conclusions**

533 Sea turtle conservation managers and scientists face difficult management choices as global
534 warming progresses, and tools are still missing to make fully informed decisions about mitigation
535 strategies. Barriers to determining neonate sex have limited our understanding of primary sex
536 ratios in this taxon, and this knowledge gap must rapidly be overcome. It is likely sea turtles have
537 evolved heritable mechanisms, whether behavioural, genetic, epigenetic or physiological, to
538 respond to climate change, but these are not currently quantified.

539 To inform future decisions, we must accept it is not sufficient to estimate sex ratios based on
540 theoretical pivotal temperatures from distant populations. We must begin to monitor sex ratios in
541 a high throughput manner to quantify the adaptive potential of these species. This will enable us
542 to assess the thermal response curves of populations and nesting aggregations, along with their
543 variation across time and space. How this variation is maintained is an important question in the
544 light of climate change and the underlying mechanisms will need to be clarified.

545 **Acknowledgements**

546 The authors would like to thank the NGOs they work with for sharing their concerns and asking
547 for converting academic discussions into practical conservation recommendations. Specifically,
548 we are grateful for our trustful interactions with Biosferal, Caretta Caretta, Foundation Biodiversity
549 Maio, Institute for Ocean (iMar), Projecto Vito Fogo, Projecto Vito Porto Novo, Project Biodiversity
550 and Turtle Foundation. This work is supported by a Natural Environment Research Council grant
551 to CE (NE/V001469/1) and Queen Mary University of London. ECL would like to thank the centre
552 for public engagement fund at QMUL for their support at the inception of this project.

553 **Data Availability Statement**

554 Data sharing not applicable to this article as no datasets were generated or analysed during the
555 current study.

556

557 **References**

558 Abreu-Grobois, F. A., Morales-Mérida, B. A., Hart, C. E., Guillon, J. M., Godfrey, M. H., Navarro,
559 E. and Girondot, M. (2020) Recent advances on the estimation of the thermal reaction norm for
560 sex ratios, *PeerJ*, 2020(3), pp. 1–28. doi: 10.7717/peerj.8451.

561

562 Ackerman, R. (1997) The nest environment and the embryonic development of sea turtles, in *The*
563 *Biology of Sea Turtles, Volume I*, pp. 83–106.

564

565 Arendt, M. D., Segars, A. L., Byrd, J. I., Boynton, J., Whitaker, J. D., Parker, L., Owens, D. W.,
566 Blanvillain, G., Quattro, J. M. and Roberts, M. A. (2012) Distributional patterns of adult male
567 loggerhead sea turtles (*Caretta caretta*) in the vicinity of Cape Canaveral, Florida, USA during
568 and after a major annual breeding aggregation, *Marine Biology*, 159, pp. 101–112. doi:
569 10.1007/s00227-011-1793-5.

570

571 Bachtrog, D., Mank, J. E., Peichel, C. L., Kirkpatrick, M., Otto, S. P., Ashman, T. L., Hahn, M. W.,
572 Kitano, J., Mayrose, I., Ming, R., Perrin, N., Ross, L., Valenzuela, N., Vamosi, J. C., Mank, J. E.,
573 Peichel, C. L., Ashman, T. L., Blackmon, H., Goldberg, E. E., Hahn, M. W., Kirkpatrick, M.,
574 Kitano, J., Mayrose, I., Ming, R., Pennell, M. W., Perrin, N., Valenzuela, N. and Vamosi, J. C.
575 (2014) Sex determination: Why so many ways of doing it?, *PLoS Biology*, 12(7), pp. 1–13. doi:
576 10.1371/journal.pbio.1001899.

577

578 Baltazar-Soares, M., Klein, J. D., Correia, S. M., Reischig, T., Taxonera, A., Roque, S. M., Dos
579 Passos, L., Durão, J., Lomba, J. P., Dinis, H., Cameron, S. J. K., Stiebens, V. A. and Eizaguirre,
580 C. (2020) Distribution of genetic diversity reveals colonization patterns and philopatry of the
581 loggerhead sea turtles across geographic scales, *Scientific Reports*, 10(1). doi: 10.1038/s41598-
582 020-74141-6.

583

584 Barnosky, A. D., Matzke, N., Tomiya, S., Wogan, G. O. U., Swartz, B., Quental, T. B., Marshall,
585 C., McGuire, J. L., Lindsey, E. L., Maguire, K. C., Mersey, B. and Ferrer, E. A. (2011) Has the
586 Earth's sixth mass extinction already arrived?, *Nature*, 471(7336), pp. 51–57. doi:

587 10.1038/nature09678.

588

589 Barske, L. A. and Capel, B. (2010) Estrogen represses SOX9 during sex determination in the red-
590 eared slider turtle *Trachemys scripta*, *Developmental Biology*, 341(1), pp. 305–314. doi:

591 10.1016/j.ydbio.2010.02.010.

592

593 Bentley, B. P., Stubbs, J. L., Whiting, S. D. and Mitchell, N. J. (2020) Variation in thermal traits
594 describing sex determination and development in Western Australian sea turtle populations,
595 *Functional Ecology*, 34(11), pp. 2302–2314. doi: 10.1111/1365-2435.13645.

596

597 Binckley, C. B., Spotila, J. R., Wilson, K. S. and Paladino, F. W. (1998) Sex determination and
598 sex ratios of pacific leatherback turtles , *Dermochelys coriacea*, *Copeia*, 2, pp. 291–300.

599

600 Bókony, V., Milne, G., Pipoly, I., Székely, T. and Liker, A. (2019) Sex ratios and bimaturism differ
601 between temperature-dependent and genetic sex-determination systems in reptiles, *BMC*
602 *Evolutionary Biology*, 19(1), pp. 1–7. doi: 10.1186/s12862-019-1386-3.

603

604 Boon, W. C. and Simpson, E. R. (2012) Neuroendocrine inherited or induced aromatase enzyme
605 deficits, *Handbook of Neuroendocrinology*. Elsevier Inc. doi: 10.1016/B978-0-12-375097-
606 6.10033-2.

607

608 Booth, D. T. and Astill, K. (2001) Temperature variation within and between nests of the green
609 sea turtle, *Chelonia mydas* (Chelonia: Cheloniidae) on Heron Island, Great Barrier Reef,
610 *Australian Journal of Zoology*, 49(1), pp. 71–84. doi: 10.1071/ZO00059.

611

612 Booth, D. T. and Freeman, C. (2006) Sand and nest temperatures and an estimate of hatchling
613 sex ratio from the Heron Island green turtle (*Chelonia mydas*) rookery, Southern Great Barrier
614 Reef, *Coral Reefs*, 25, pp. 629–633. doi: 10.1007/s00338-006-0135-4.

615

616 Bowden, R. M., Ewert, M. A., Freedberg, S. and Nelson, C. E. (2002) Maternally derived yolk
617 hormones vary in follicles of the painted turtle, *Chrysemys picta*, *Journal of Experimental*
618 *Zoology*, 293(1), pp. 67–72. doi: 10.1002/jez.10094.

619

620 Bowden, R. M., Ewert, M. A. and Nelson, C. E. (2000) Environmental sex determination in a
621 reptile varies seasonally and with yolk hormones, *Proceedings of the Royal Society B: Biological*

622 Sciences, 267(1454), pp. 1745–1749. doi: 10.1098/rspb.2000.1205.

623

624 Bowden, R. M. and Paitz, R. T. (2018) Temperature fluctuations and maternal estrogens as

625 critical factors for understanding temperature-dependent sex determination in nature, *Journal of*

626 *Experimental Biology part A: Ecological and Integrative Physiology*, pp. 1–8. doi:

627 10.1002/jez.2183.

628

629 Bowen, B. W. and Karl, S. A. (2007) Population genetics and phylogeography of sea turtles,

630 *Molecular Ecology*, 16, pp. 4886–4907. doi: 10.1111/j.1365-294X.2007.03542.x.

631

632 Broderick, A. C., Godley, B. J. and Hays, G. C. (2001) Trophic status drives interannual variability

633 in nesting numbers of marine turtles, *Proceedings of the Royal Society B: Biological Sciences*,

634 268(1475), pp. 1481–1487. doi: 10.1098/rspb.2001.1695.

635

636 Bull, J. J. and Vogt, R. C. (1979) Temperature-dependent sex determination in turtles, *Science*,

637 206(4423), pp. 1186–1188.

638

639 Bulun, S. E., Takayama, K., Suzuki, T., Sasano, H., Yilmaz, B. and Sebastian, S. (2004)

640 Organization of the human Aromatase P450 (CYP19) gene, *Seminars in Reproductive Medicine*,

641 22(1), pp. 5–9.

642

643 Carter, A. W., Bowden, R. M. and Paitz, R. T. (2017) Seasonal shifts in sex ratios are mediated

644 by maternal effects and fluctuating incubation temperatures, *Functional Ecology*, 31, pp. 876–

645 884. doi: 10.1111/1365-2435.12801.

646

647 Charnier, M. (1966) Action of temperature on the sex ratio in the *Agama agama* (Agamidae,

648 Lacertilia) embryo, *C R Seances Soc Biol Fil.*, 160(3), pp. 620–622.

649

650 Charnov, E. . and Bull, J. . (1977) When is sex environmentally determined?, *Nature*, 266, pp.

651 828–830.

652

653 Chevin, L. M., Lande, R. and Mace, G. M. (2010) Adaptation, plasticity, and extinction in a

654 changing environment: Towards a predictive theory, *PLoS Biology*, 8(4). doi:

655 10.1371/journal.pbio.1000357.

656

657 Conley, A. and Hinshelwood, M. (2001) Mammalian aromatases, *Reproduction*, 121(5), pp. 685–
 658 695. doi: 10.1530/rep.0.1210685.

659

660 Cree, A., Thompson, M. B. and Daugherty, C. H. (1995) Tuatara sex determination, *Nature*, 375,
 661 p. 543.

662

663 Crews, D., Bull, J. J. and Wibbels, T. (1991) Estrogen and sex reversal in turtles: A dose-
 664 dependent phenomenon, *General and Comparative Endocrinology*, 81(3), pp. 357–364. doi:
 665 10.1016/0016-6480(91)90162-Y.

666

667 Crews, D., Wibbels, T. and Gutzke, W. H. N. (1989) Action of sex steroid hormones on
 668 temperature-induced sex determination in the snapping turtle (*Chelydra serpentina*), *General and*
 669 *Comparative Endocrinology*, 76(1), pp. 159–166. doi: 10.1016/0016-6480(89)90042-7.

670

671 Deeming, D. C., Ferguson, M. W. J., Mittwoch, U., Wolf, U., Dorizzi, M., Zaborski, P. and Sharma,
 672 H. (1988) Environmental regulation of sex determination in reptiles (and discussion),
 673 *Philosophical Transactions of the Royal Society B: Biological Sciences*, 322(1208), pp. 19–39.

674

675 DeGregorio, B. A. and Williard, A. S. (2011) Incubation temperatures and metabolic heating of
 676 relocated and in situ loggerhead sea turtle (*Caretta caretta*) nests at a northern rookery,
 677 *Chelonian Conservation and Biology*, 10(1), pp. 54–61. doi: 10.2744/CCB-0880.1.

678

679 Dellert, L. J., O’Neil, D. and Cassill, D. L. (2014) Effects of beach renourishment and clutch
 680 relocation on the success of the loggerhead sea turtle (*Caretta caretta*) eggs and hatchlings,
 681 *Journal of Herpetology*, 48(2), pp. 186–187. doi: 10.1670/12-135.

682

683 Deveson, I. W., Holleley, C. E., Blackburn, J., Marshall Graves, J. A., Mattick, J. S., Waters, P. D.
 684 and Georges, A. (2017) Differential intron retention in Jumonji chromatin modifier genes is
 685 implicated in reptile temperature-dependent sex determination, *Science Advances*, 3(6), pp. 1–8.
 686 doi: 10.1126/sciadv.1700731.

687

688 Díaz-Hernández, V., Marmolejo-Valencia, A. and Merchant-Larios, H. (2015) Exogenous
 689 estradiol alters gonadal growth and timing of temperature sex determination in gonads of sea
 690 turtle, *Developmental Biology*, 408(1), pp. 79–89. doi: 10.1016/j.ydbio.2015.05.022.

691

692 Díaz-Hernández, V., Vázquez-Gómez, A., Marmolejo-Valencia, A., Montaña, L. M. and Merchant-
 693 Larios, H. (2017) 17β -Estradiol modulates cell proliferation of medullary cords during ovarian
 694 differentiation of the *Lepidochelys olivacea* sea turtle, *Developmental Biology*, pp. 1–9. doi:
 695 10.1016/j.ydbio.2017.09.008.
 696
 697 Dodd, K. L. and Wibbels, T. (2008) Estrogen inhibits caudal progression but stimulates
 698 proliferation of developing müllerian ducts in a turtle with temperature-dependent sex
 699 determination, *Comparative Biochemistry and Physiology - A Molecular and Integrative*
 700 *Physiology*, 150(3), pp. 315–319. doi: 10.1016/j.cbpa.2008.04.002.
 701
 702 Eberhart-Phillips, L. J., Küpper, C., Miller, T. E. X., Cruz-López, M., Maher, K. H., dos Remedios,
 703 N., Stoffel, M. A., Hoffman, J. I., Krüger, O. and Székely, T. (2017) Sex-specific early survival
 704 drives adult sex ratio bias in snowy plovers and impacts mating system and population growth,
 705 *Proceedings of the National Academy of Sciences of the United States of America*, 114(27), pp.
 706 E5474 - E5481, doi: 10.1073/pnas.1620043114
 707
 708
 709 Edwards, A. M., Cameron, E. Z., Wapstra, E. and McEvoy, J. (2019) Maternal effects obscure
 710 condition-dependent sex allocation in changing environments, *Royal Society Open Science*, 6(4).
 711 doi: 10.1098/rsos.181885.
 712
 713 Eizaguirre, C. and Baltazar-Soares, M. (2014) Evolutionary conservation-evaluating the adaptive
 714 potential of species, *Evolutionary Applications*, 7(9), pp. 963–967. doi: 10.1111/eva.12227.
 715
 716 Elf, P. K. (2003) Yolk steroid hormones and sex determination in reptiles with TSD, *General and*
 717 *Comparative Endocrinology*, 132(3), pp. 349–355. doi: 10.1016/S0016-6480(03)00098-4.
 718
 719 Esteban, N., Laloë, J. O., Kiggen, F. S. P. L., Ubels, S. M., Becking, L. E., Meesters, E. H.,
 720 Berkel, J., Hays, G. C. and Christianen, M. J. A. (2018) Optimism for mitigation of climate
 721 warming impacts for sea turtles through nest shading and relocation, *Scientific Reports*, 8(1), pp.
 722 1–8. doi: 10.1038/s41598-018-35821-6.
 723
 724 Esteban, N., Laloë, J. O., Mortimer, J. A., Guzman, A. N. and Hays, G. C. (2016) Male hatchling
 725 production in sea turtles from one of the world's largest marine protected areas, the Chagos

726 Archipelago, *Scientific Reports* 6:20339, pp. 1–8. doi: 10.1038/srep20339.

727

728 Ewert, M. A., Lang, J. W. and Nelson, C. E. (2005) Geographic variation in the pattern of

729 temperature-dependent sex determination in the American snapping turtle (*Chelydra serpentina*),

730 *Journal of Zoology*, 265(1), pp. 81–95. doi: 10.1017/S0952836904006120.

731

732 Ferguson, M. W. J. and Joanen, T. (1983) Temperature-dependent sex determination in *Alligator*

733 *mississippiensis*, *Journal of Zoology*, 200(2), pp. 143–177. doi: 10.1111/j.1469-

734 7998.1983.tb05781.x.

735

736 Fisher, C. R., Graves, K. H., Parlow, A. F. and Simpson, E. R. (1998) Characterization of mice

737 deficient in aromatase (ArKO) because of targeted disruption of the cyp19 gene, *Proceedings of*

738 *the National Academy of Sciences of the United States of America*, 95, pp. 6965–6970. doi:

739 10.1073/pnas.95.12.6965.

740

741 Fisher, R. A. (1930) *The genetical theory of natural selection*. Clarendon Press, Oxford.

742

743 Fossette, S., Witt, M. J., Miller, P., Nalovic, M. A., Albareda, D., Almeida, A. P., Broderick, A. C.,

744 Chacon-Chaverri, D., Coyne, M. S., Domingo, A., Eckert, S., Evans, D., Fallabrino, A., Ferraroli,

745 S., Formia, A., Giffoni, B., Hays, G. C., Hughes, G., Kelle, L., Leslie, A., Lopez-Mendilaharsu, M.,

746 Luschi, P., Prosdocimi, L., Rodriguez-Heredia, S., Turny, A., Verhage, S. and Godley, B. J.

747 (2014) Pan-Atlantic analysis of the overlap of a highly migratory species , the leatherback turtle ,

748 with pelagic longline fisheries, *Proceedings of The Royal Society B*, 281, p. 20133065.

749

750 Frankham, R. (2005) Genetics and extinction, *Biological Conservation*, 126, pp. 131–140. doi:

751 10.1016/j.biocon.2005.05.002.

752

753 Fuentes, M. M. P. B., Fish, M. R. and Maynard, J. A. (2012) Management strategies to mitigate

754 the impacts of climate change on sea turtle's terrestrial reproductive phase, *Mitigation and*

755 *Adaptation Strategies for Global Change*, 17(1), pp. 51–63. doi: 10.1007/s11027-011-9308-8.

756

757 Fuentes, M. M. P. B., Monsinjon, J., Lopez, M., Lara, P., Santos, A., dei Marcovaldi, M. A. G. and

758 Girondot, M. (2017) Sex ratio estimates for species with temperature-dependent sex

759 determination differ according to the proxy used, *Ecological Modelling*, 365, pp. 55–67. doi:

760 10.1016/j.ecolmodel.2017.09.022.

761

762 Ge, C., Ye, J., Weber, C., Sun, W., Zhang, H., Zhou, Y., Cai, C., Qian, G. and Capel, B. (2018)
763 The histone demethylase KDM6B regulates temperature-dependent sex determination in a turtle
764 species, *Science*, 360(6389), pp. 645–648. doi: 10.1126/science.aap8328.

765

766 Ge, C., Ye, J., Zhang, H., Zhang, Y., Sun, W., Sang, Y., Capel, B. and Qian, G. (2017) Dmrt1
767 induces the male pathway in a turtle species with temperature-dependent sex determination,
768 *Development (Cambridge)*, 144(12), pp. 2222–2233. doi: 10.1242/dev.152033.

769

770 Georges, A., Young, J. E. and Doody, J. S. (2005) Modelling Development of Reptile Embryos
771 under Fluctuating Temperature Regimes, 78(1), pp. 18–30.

772

773 Gienapp, P., Teplitsky, C., Alho, J. S., Mills, J. A. and Merilä, J. (2008) Climate change and
774 evolution: Disentangling environmental and genetic responses, *Molecular Ecology*, 17(1), pp.
775 167–178. doi: 10.1111/j.1365-294X.2007.03413.x.

776

777 Girondot, M. (1999) Statistical description of temperature-dependent sex determination using
778 maximum likelihood, *Evolutionary Ecology Research*, 1(4), pp. 479–486.

779

780 Girondot, M. and Pieau, C. (1999) A fifth hypothesis for the evolution of TSD in reptiles, *Trends in*
781 *Ecology & Evolution*, 14(9), pp. 359–360. doi: 10.1016/s0169-5347(99)01681-x.

782

783 Glen, F. and Mrosovsky, N. (2004) Antigua revisited: The impact of climate change on sand and
784 nest temperatures at a hawksbill turtle (*Eretmochelys imbricata*) nesting beach, *Global Change*
785 *Biology*, 10(12), pp. 2036–2045. doi: 10.1111/j.1529-8817.2003.00865.x.

786

787 Godfrey, M. H., D'Amato, A. F., Marcovaldi, M. Â. and Mrosovsky, N. (1999) Pivotal temperature
788 and predicted sex ratios for hatchling hawksbill turtles from Brazil, *Canadian Journal of Zoology*,
789 77(9), pp. 1465–1473. doi: 10.1139/z99-117.

790

791 Godfrey, M. H., Mrosovsky, N. and Barreto, R. (1996) Estimating past and present sex ratios of
792 sea turtles in Suriname, *Canadian Journal of Zoology*, 74, pp. 267–277. doi: 10.1139/z96-033.

793

794 González, E. J., Martínez-López, M., Morales-Garduza, M. A., García-Morales, R., Charruau, P.
795 and Gallardo-Cruz, J. A. (2019) The sex-determination pattern in crocodilians: A systematic

796 review of three decades of research, *Journal of Animal Ecology*, 88(9), pp. 1417–1427. doi:
 797 10.1111/1365-2656.13037.
 798

799 Grant, V. J. (2007) Could maternal testosterone levels govern mammalian sex ratio deviations?,
 800 *Journal of Theoretical Biology*, 246(4), pp. 708–719. doi: 10.1016/j.jtbi.2007.02.005.
 801

802 Gross, T. S., Crain, D. A., Bjorndal, K. A., Bolten, A. B. and Carthy, R. R. (1995) Identification of
 803 sex in hatchling loggerhead turtles (*Caretta caretta*) by analysis of steroid concentrations in
 804 chorioallantoic/amniotic fluid., *General and comparative endocrinology*, pp. 204–210. doi:
 805 10.1006/gcen.1995.1103.
 806

807 Guiguen, Y., Fostier, A., Piferrer, F. and Chang, C. F. (2010) Ovarian aromatase and estrogens:
 808 A pivotal role for gonadal sex differentiation and sex change in fish, *General and Comparative*
 809 *Endocrinology*, 165(3), pp. 352–366. doi: 10.1016/j.ygcen.2009.03.002.
 810

811 Hamann, M., Godfrey, M. H., Seminoff, J. A., Arthur, K., Barata, P. C. R., Bjorndal, K. A., Bolten,
 812 A. B., Broderick, A. C., Campbell, L. M., Carreras, C., Casale, P., Chaloupka, M., Chan, S. K. F.,
 813 Coyne, M. S., Crowder, L. B., Diez, C. E., Dutton, P. H., Epperly, S. P., Fitz Simmons, N. N.,
 814 Formia, A., Girondot, M., Hays, G. C., Cheng, I. J., Kaska, Y., Lewison, R., Mortimer, J. A.,
 815 Nichols, W. J., Reina, R. D., Shanker, K., Spotila, J. R., Tomás, J., Wallace, B. P., Work, T. M.,
 816 Zbinden, J. and Godley, B. J. (2010) Global research priorities for sea turtles: Informing
 817 management and conservation in the 21st century, *Endangered Species Research*, 11(3), pp.
 818 245–269. doi: 10.3354/esr00279.
 819

820 Hanson, J. A., Wibbels, T. and Martin, R. E. (1998) Predicted female bias in sex ratios of
 821 hatchling loggerhead sea turtles from a Florida nesting beach, *Canadian Journal of Zoology*, 76,
 822 pp. 1850–1861. doi: 10.1139/z98-118.
 823

824 Hawkes, L. A., Broderick, A. C., Coyne, M. S., Godfrey, M. H. and Godley, B. J. (2007) ‘Only
 825 some like it hot - Quantifying the environmental niche of the loggerhead sea turtle’, *Diversity and*
 826 *Distributions*, 13(4), pp. 447–457. doi: 10.1111/j.1472-4642.2007.00354.x.
 827

828 Hawkes, L. A., Broderick, A. C., Godfrey, M. H. and Godley, B. J. (2007) Investigating the
 829 potential impacts of climate change on a marine turtle population, *Global Change Biology*, 13(5),
 830 pp. 923–932. doi: 10.1111/j.1365-2486.2007.01320.x.

831

832 Hawkes, L. A., Broderick, A., Godfrey, M. and Godley, B. (2009) Climate change and marine
833 turtles, *Endangered Species Research*, 7(May), pp. 137–154. doi: 10.3354/esr00198.

834

835 Hays, G. C., Ashworth, J. S., Barnsley, M. J., Broderick, A. C., Emery, D. R., Godley, B. J.,
836 Henwood, A. and Jones, E. L. (2001) The importance of sand albedo for the thermal conditions
837 on sea turtle nesting beaches, *Oikos*, 93, pp. 87–94.

838

839 Hays, G. C., Fossette, S., Katselidis, K. A., Schofield, G. and Gravenor, M. B. (2010) Breeding
840 periodicity for male sea turtles, operational sex ratios, and implications in the face of climate
841 change, *Conservation Biology*, 24(6), pp. 1636–1643. doi: 10.1111/j.1523-1739.2010.01531.x.

842

843 Hays, G. C., Mazaris, A. D. and Schofield, G. (2014) Different male vs. female breeding
844 periodicity helps mitigate offspring sex ratio skews in sea turtles, *Frontiers in Marine Science*,
845 1(September), pp. 1–9. doi: 10.3389/fmars.2014.00043.

846

847 Hedrick, P. W. and Kalinowski, S. T. (2000) Inbreeding depression in conservation biology,
848 *Annual Review of Ecology, Evolution, and Systematics*, 31, pp. 139–162.

849

850 Hoffmann, A. and Sgrò, C. (2011) Climate change and evolutionary adaptation., *Nature*,
851 470(7335), pp. 479–485. doi: 10.1038/nature09670.

852

853 Von Holle, B., Irish, J. L., Spivy, A., Weishampel, J. F., Meylan, A., Godfrey, M. H., Dodd, M.,
854 Schweitzer, S. H., Keyes, T., Sanders, F., Chaplin, M. K. and Taylor, N. R. (2019) Effects of
855 future sea level rise on coastal habitat, *Journal of Wildlife Management*, 83(3), pp. 694–704. doi:
856 10.1002/jwmg.21633.

857

858 Holleley, C. E., O'Meally, D., Sarre, S. D., Marshall Graves, J. A., Ezaz, T., Matsubara, K., Azad,
859 B., Zhang, X. and Georges, A. (2015) Sex reversal triggers the rapid transition from genetic to
860 temperature-dependent sex, *Nature*, 523(7558), pp. 79–82. doi: 10.1038/nature14574.

861

862 Holleley, C. E., Sarre, S. D., O'Meally, D. and Georges, A. (2016) Sex reversal in reptiles:
863 reproductive oddity or powerful driver of evolutionary change?, *Sexual Development*, 10(5–6), pp.
864 279–287. doi: 10.1159/000450972.

865

866 Houghton, J. D. R., Myers, A. E., Lloyd, C., King, R. S., Isaacs, C. and Hays, G. C. (2007)
 867 Protracted rainfall decreases temperature within leatherback turtle (*Dermochelys coriacea*)
 868 clutches in Grenada, West Indies: Ecological implications for a species displaying temperature
 869 dependent sex determination, *Journal of Experimental Marine Biology and Ecology*, 345(1), pp.
 870 71–77. doi: 10.1016/j.jembe.2007.02.001.

871

872 Howard, R., Bell, I. and Pike, D. A. (2015) Tropical flatback turtle (*Natator depressus*) embryos
 873 are resilient to the heat of climate change, *Journal of Experimental Biology*, 218(20), pp. 3330–
 874 3335. doi: 10.1242/jeb.118778.

875

876 Janes, D. E., Bermudez, D., Guillette, L. J. and Wayne, M. L. (2007) Estrogens Induced Male
 877 Production at a Female-Producing Temperature in a Reptile (Leopard Gecko, *Eublepharis*
 878 *Macularius*) with Temperature-dependent Sex Determination, *Journal of Herpetology*, 41(1), pp.
 879 9–15. doi: 10.1670/0022-1511(2007)41[9:EIMPAA]2.0.CO;2.

880

881 Janzen, F. J. and Krenz, J. D. (2004) Phylogenetics: Which was first, TSD or GSD?, in
 882 *Temperature-Dependent Sex Determination in Vertebrates*, pp. 121–130.

883

884 Janzen, F. J. and Paukstis, G. L. (1991) Environmental Sex Determination in Reptiles: Ecology,
 885 Evolution and Experimental Design, *The Quarterly Review of Biology*, 66(2), pp. 149–179.

886

887 Janzen, F. J. and Phillips, P. C. (2006) Exploring the evolution of environmental sex
 888 determination, especially in reptiles, *Journal of Evolutionary Biology*, 19(6), pp. 1775–1784. doi:
 889 10.1111/j.1420-9101.2006.01138.x.

890

891 Jeyasuria, P. and Place, A. R. (1998) Embryonic brain-gonadal axis in temperature-dependent
 892 sex determination of reptiles: A role for P450 aromatase (CYP19), *Journal of Experimental*
 893 *Zoology*, 281, pp. 428–449. doi: 10.1002/(SICI)1097-010X(19980801)281:5<428::AID-
 894 JEZ8>3.0.CO;2-Q.

895

896 Jribi, I. and Bradai, M. N. (2014) Sex ratio estimations of loggerhead sea turtle hatchlings at
 897 Kuriat Islands, Tunisia: Can minor nesting sites contribute to compensate globally female-biased
 898 sex ratio?, *Scientific World Journal*, 2014. doi: 10.1155/2014/419410.

899

900 Kaska, Y., Başkale, E., Urhan, R., Katılmış, Y., Gidiş, M., Sarı, F., Sözbilen, D., Canbolat, A. F.,

901 Yılmaz, F., Barlas, M., Özdemir, N. and Özkul, M. (2013) Natural and anthropogenic factors
 902 affecting the nest-site selection of loggerhead turtles, *Caretta caretta*, on Dalaman-Sarigerme
 903 beach in South- west Turkey, *Zoology in the Middle East*, 50(1), pp. 47–58. doi:
 904 10.1080/09397140.2010.10638411.

905

906 Kaska, Y., Downie, R., Tippet, R. and Furness, R. W. (1998) Natural temperature regimes for
 907 loggerhead and green turtle nests in the eastern Mediterranean, *Canadian journal of zoology*, 76,
 908 pp. 723–729.

909

910 Ko, C. I., Chesnel, A., Mazerbourg, S., Kuntz, S., Flament, S. and Chardard, D. (2008) Female-
 911 enriched expression of ER α during gonad differentiation of the urodele amphibian *Pleurodeles*
 912 *waltl*, *General and Comparative Endocrinology*, 156(2), pp. 234–245. doi:
 913 10.1016/j.ygcen.2008.01.012.

914

915 Laloë, J.-O., Cozens, J., Renom, B., Taxonera, A., Hays, G. C., Vicente, S., Nicolau, S., Boa
 916 Vista, S., Santiago, M. and Brava, F. (2014) Climate change induced extinction does not seem
 917 imminent at an important sea turtle rookery, *Nature Climate Change*, 4, pp. 1–7. doi:
 918 10.1038/NCLIMATE2236.

919

920 Lande, R. (2009) Adaptation to an extraordinary environment by evolution of phenotypic plasticity
 921 and genetic assimilation, *Journal of Evolutionary Biology*, 22, pp. 1435–1446. doi:
 922 10.1111/j.1420-9101.2009.01754.x.

923

924 Lange, I. G., Hartel, A. and Meyer, H. H. D. (2002) Evolution of oestrogen functions in
 925 vertebrates, *Journal of Steroid Biochemistry and Molecular Biology*, 83(1–5), pp. 219–226. doi:
 926 10.1016/S0960-0760(02)00225-X.

927

928 Lee, P. L. M. and Hays, G. C. (2004) Polyandry in a marine turtle: Females make the best of a
 929 bad job, *Proceedings of the National Academy of Sciences*, 101(17), pp. 6530–6535. doi:
 930 10.1073/pnas.0307982101.

931

932 Lee, P. L. M., Luschi, P. and Hays, G. C. (2007) Detecting female precise natal philopatry in
 933 green turtles using assignment methods, *Molecular Ecology*, 16, pp. 61–74. doi: 10.1111/j.1365-
 934 294X.2006.03115.x.

935

936 Levasseur, K. E., Stapleton, S. P. and Quattro, J. M. (2020) Precise natal homing and an
 937 estimate of age at sexual maturity in hawksbill turtles, *Animal Conservation*. doi:
 938 10.1111/acv.12657.
 939
 940 Lockley, E. C., Reischig, T. and Eizaguirre, C. (2020) Maternally derived sex steroid hormones
 941 impact sex ratios of loggerhead sea turtles, *bioRxiv*, p. 2020.01.10.901520. doi:
 942 10.1101/2020.01.10.901520.
 943
 944 Lolavar, A. and Wyneken, J. (2015) Effect of rainfall on loggerhead turtle nest temperatures, sand
 945 temperatures and hatchling sex, *Endangered Species Research*, 28(3), pp. 235–247. doi:
 946 10.3354/esr00684.
 947
 948 Lolavar, A. and Wyneken, J. (2017) Experimental assessment of the effects of moisture on
 949 loggerhead sea turtle hatchling sex ratios, *Zoology*, 123, pp. 64–70. doi:
 950 10.1016/j.zool.2017.06.007.
 951
 952 Lolavar, A. and Wyneken, J. (2020) The impact of sand moisture on the temperature-sex ratio
 953 responses of developing loggerhead (*Caretta caretta*) sea turtles, *Zoology*, 138. doi:
 954 10.1016/j.zool.2019.125739.
 955
 956 Marcovaldi, M., Godfrey, M. H. and Mrosovsky, N. (1997) Estimating sex ratios of loggerhead
 957 turtles in Brazil from pivotal incubation durations, *Canadian Journal of Zoology*, 75, pp. 755–770.
 958
 959 Martínez-Juárez, A. and Moreno-Mendoza, N. (2019) Mechanisms related to sexual
 960 determination by temperature in reptiles, *Journal of Thermal Biology*. Elsevier Ltd. doi:
 961 10.1016/j.jtherbio.2019.102400.
 962
 963 Massey, M. D., Holt, S. M., Brooks, R. J. and Rollinson, N. (2019) Measurement and modelling of
 964 primary sex ratios for species with temperature-dependent sex determination, *Journal of*
 965 *Experimental Biology*, 222(1). doi: 10.1242/jeb.190215.
 966
 967 Matsumoto, Y., Hannigan, B. and Crews, D. (2016) Temperature Shift Alters DNA Methylation
 968 and Histone Modification Patterns in Gonadal Aromatase (cyp19a1) Gene in Species with
 969 Temperature-Dependent Sex Determination, *Plos One*, 11(11), p. e0167362. doi:
 970 10.1371/journal.pone.0167362.

971

972 Matsuzawa, Y., Sato, K., Sakamoto, W. and Bjorndal, K. A. (2002) Seasonal fluctuations in sand
973 temperature: Effects on the incubation period and mortality of loggerhead sea turtle (*Caretta*
974 *caretta*) pre-emergent hatchlings in Minabe, Japan, *Marine Biology*, 140(3), pp. 639–646. doi:
975 10.1007/s00227-001-0724-2.

976

977 Maulany, R. I., Booth, D. T. and Baxter, G. S. (2012) The effect of incubation temperature on
978 hatchling quality in the olive ridley turtle, *Lepidochelys olivacea*, from Alas Purwo National Park,
979 East Java, Indonesia: Implications for hatchery management, *Marine Biology*, 159(12), pp. 2651–
980 2661. doi: 10.1007/s00227-012-2022-6.

981

982 Mazaris, A. D., Kallimanis, A. S., Pantis, J. D. and Hays, G. C. (2013) Phenological response of
983 sea turtles to environmental variation across a species' northern range., *Proceedings. Biological*
984 *sciences / The Royal Society*, 280(1751), p. 20122397. doi: 10.1098/rspb.2012.2397.

985

986 McCoy, C. J., Vogt, R. C. and Censky, E. J. (1983) Temperature-controlled sex determination in
987 the sea turtle *Lepidochelys olivacea*, *Journal of Herpetology*, 17(4), pp. 404–406.

988

989 McGaugh, S. E., Bowden, R. M., Kuo, C. H. and Janzen, F. J. (2011) Field-measured heritability
990 of the threshold for sex determination in a turtle with temperature-dependent sex determination,
991 *Evolutionary Ecology Research*, 13(1), pp. 75–90.

992

993 McMahon, C. R. and Hays, G. C. (2006) Thermal niche, large-scale movements and implications
994 of climate change for a critically endangered marine vertebrate, *Global Change Biology*, 12(7),
995 pp. 1330–1338. doi: 10.1111/j.1365-2486.2006.01174.x.

996

997 Merchant-Larios, H., Ruiz-Ramirez, S., Moreno-Mendoza, N. and Marmolejo-Valencia, a (1997)
998 Correlation among thermosensitive period, estradiol response, and gonad differentiation in the
999 sea turtle *Lepidochelys olivacea*., *General and comparative endocrinology*, 107(3), pp. 373–385.
1000 doi: 10.1006/gcen.1997.6946.

1001

1002 Merilä, J. and Hendry, A. P. (2014) Climate change, adaptation, and phenotypic plasticity: The
1003 problem and the evidence, *Evolutionary Applications*, 7(1), pp. 1–14. doi: 10.1111/eva.12137.

1004

1005 Meyers, L. A. and Bull, J. J. (2002) Fighting change with change: adaptive variation in an

1006 uncertain world, *Trends Ecol. Evol.*, 17(12), pp. 551–557.

1007

1008 Meylan, A. B., Bowen, B. W. and Avise, J. C. (1990) A Genetic Test of the Natal Homing Versus

1009 Social Facilitation Models for Green Turtle Migration, *Science*, 248, pp. 724–728.

1010

1011 Mitchell, N. J. and Janzen, F. J. (2010) Temperature-Dependent sex determination and

1012 contemporary climate change, *Sexual Development*, 4(1–2), pp. 129–140. doi:

1013 10.1159/000282494.

1014

1015 Monsinjon, J. R., Wyneken, J., Rusenko, K., López-Mendilaharsu, M., Lara, P., Santos, A., dei

1016 Marcovaldi, M. A. G., Fuentes, M. M. P. B., Kaska, Y., Tucek, J., Nel, R., Williams, K. L., LeBlanc,

1017 A. M., Rostal, D., Guillon, J. M. and Girondot, M. (2019) The climatic debt of loggerhead sea

1018 turtle populations in a warming world,

1019 *Ecological Indicators*, 107(May), p. 105657. doi: 10.1016/j.ecolind.2019.105657.

1020

1021 Morjan, A. C. L. and Janzen, F. J. (2003) Nest temperature is not related to egg size in a turtle

1022 with temperature-dependent sex determination, *Copeia*, (2), pp. 366–372.

1023

1024 Morreale, S. J., Ruiz, G. J., Spotila, J. R. and Standora, E. A. (1982) Temperature-dependent sex

1025 determination: Current practices threaten conservation of sea turtles. *Science*, 216, pp. 1245–

1026 1247.

1027

1028 Morrish, B. C. and Sinclair, A. H. (2002) Vertebrate sex determination: Many means to an end,

1029 *Reproduction*, 124(4), pp. 447–457. doi: 10.1530/rep.0.1240447.

1030

1031 Mortimer, J. A. and Donnelly, M. (2008) *Eretmochelys imbricata*, *The IUCN Red List of*

1032 *Threatened Species*, p. e.T8005A12881238.

1033

1034 Mrosovsky, N. (1988) Pivotal temperatures for loggerhead turtles (*Caretta caretta*) from northern

1035 and southern nesting beaches, *Canadian Journal of Zoology*, 66(3), pp. 661–669. doi:

1036 10.1139/z88-098.

1037

1038 Mrosovsky, N. (2006) Distorting gene pools by conservation: Assessing the case of doomed turtle

1039 eggs, *Environmental Management*, 38(4), p. 703. doi: 10.1007/s00267-006-8348-4.

1040

1041 Mrosovsky, N. (2008) Against oversimplifying the issues on relocating turtle eggs. *Environmental*
 1042 *Management*, 41(4), pp. 465–467. doi: 10.1007/s00267-007-9044-8.

1043

1044 Mrosovsky, N., Bass, A., Corliss, L. A., Richardson, J. I. and Richardson, T. H. (1992) Pivotal and
 1045 beach temperatures for hawksbill turtles nesting in Antigua, *Canadian Journal of Zoology*, 70(10),
 1046 pp. 1920–1925. doi: 10.1139/z92-261.

1047

1048 Mrosovsky, N., Kamel, S., Rees, A. F. and Margaritoulis, D. (2002) Pivotal temperature for
 1049 loggerhead turtles (*Caretta caretta*) from Kyparissia Bay, Greece, *Canadian Journal of Zoology*,
 1050 80(12), pp. 2118–2124. doi: 10.1139/z02-204.

1051

1052 Mrosovsky, N. and Pieau, C. (1991) Transitional range of temperature, pivotal temperatures and
 1053 thermosensitive stages for sex determination in reptiles, *Amphibia Reptilia*, 12(2), pp. 169–179.
 1054 doi: 10.1163/156853891X00149.

1055

1056 Muralidhar, P. and Veller, C. (2018) Sexual antagonism and the instability of environmental sex
 1057 determination, *Nature Ecology & Evolution*. doi: 10.1038/s41559-017-0427-9.

1058

1059 Nakabayashi, O., Kikuchi, H., Kikuchi, T. and Mizuno, S. (1998) Differential expression of genes
 1060 for aromatase and estrogen receptor during the gonadal development in chicken embryos,
 1061 *Journal of Molecular Endocrinology*, 20(2), pp. 193–202. doi: 10.1677/jme.0.0200193.

1062

1063 Navarro-Martín, L., Viñas, J., Ribas, L., Díaz, N., Gutiérrez, A., Di Croce, L. and Piferrer, F.
 1064 (2011) DNA methylation of the gonadal aromatase (cyp19a) promoter is involved in temperature-
 1065 dependent sex ratio shifts in the European sea bass, *PLoS Genetics*, 7(12), p. e1002447. doi:
 1066 10.1371/journal.pgen.1002447.

1067

1068 Nelson, D. R., Goldstone, J. V. and Stegeman, J. J. (2013) The cytochrome P450 genesis locus:
 1069 the origin and evolution of animal cytochrome P450s, *Philos Trans R Soc Lond B Biol Sci*, 368,
 1070 20120474, doi: 10.101098/rtsb.2012.0474

1071

1072 Neukom, R., Steiger, N., Gómez-Navarro, J. J., Wang, J. and Werner, J. P. (2019) No evidence
 1073 for globally coherent warm and cold periods over the preindustrial Common Era, *Nature*, 571, pp.
 1074 550–554. doi: 10.1038/s41586-019-1401-2.

1075

- 1076 Öz, M., Erdoğan, A., Kaska, Y., Düşen, S., Aslan, A., Sert, H., Yavuz, M. and Tunç, M. R. (2004)
1077 Nest temperatures and sex-ratio estimates of loggerhead turtles at Patara beach on the
1078 southwestern coast of Turkey, *Canadian Journal of Zoology*, 82, pp. 94–101. doi: 10.1139/z03-
1079 200.
- 1080
- 1081 Patel, S. H., Morreale, S. J., Saba, V. S., Panagopoulou, A., Margaritoulis, D. and Spotila, J. R.
1082 (2016) Climate impacts on sea turtle breeding phenology in Greece and associated foraging
1083 habitats in the wider Mediterranean region, *Plos One*, 11(6), p. e0157170. doi:
1084 10.1371/journal.pone.0157170.
- 1085
- 1086 Patrício, A. R., Marques, A., Barbosa, C., Broderick, A. C., Godley, B. J., Hawkes, L. A., Rebelo,
1087 R., Regalla, A. and Catry, P. (2017) Balanced primary sex ratios and resilience to climate change
1088 in a major sea turtle population, *Marine Ecology Progress Series*, 577(August), pp. 189–203. doi:
1089 10.3354/meps12242.
- 1090
- 1091 Pen, I., Uller, T., Feldmeyer, B., Harts, A., While, G. M. and Wapstra, E. (2010) Climate-driven
1092 population divergence in sex-determining systems, *Nature*, 468(7322), pp. 436–438. doi:
1093 10.1038/nature09512.
- 1094
- 1095 Pfaller, J. B., Limpus, C. J. and Bjorndal, K. A. (2009) Nest-site selection in individual loggerhead
1096 turtles and consequences for doomed-egg relocation, *Conservation Biology*, 23(1), pp. 72–80.
1097 doi: 10.1111/j.1523-1739.2008.01055.x.
- 1098
- 1099 Pike, D. A. (2008) The benefits of nest relocation extend far beyond recruitment: A rejoinder to
1100 Mrosovsky, *Environmental Management*, 41(4), pp. 461–464. doi: 10.1007/s00267-006-0434-0.
- 1101
- 1102 Pokorná, M. J. and Kratochvil, L. (2009) Phylogeny of sex-determining mechanisms in squamate
1103 reptiles : are sex chromosomes an evolutionary trap ?, *Zoological Journal of the Linnean Society*,
1104 156(May), pp. 168–183. doi: 10.1111/j.1096-3642.2008.00481.x.
- 1105
- 1106 Quinn, A. E., Georges, A., Sarre, S. D., Guarino, F., Ezaz, T. and Graves, J. A. M. (2007)
1107 Temperature sex reversal implies sex gene dosage in a reptile, *Science*, p. 411. doi:
1108 10.1126/science.1135925.
- 1109
- 1110 Quinn, A. E., Sarre, S. D., Ezaz, T., Graves, J. A. M. and Georges, A. (2011) Evolutionary

1111 transitions between mechanisms of sex determination in vertebrates, *Biology Letters*, 7(3), pp.
 1112 443–448. doi: 10.1098/rsbl.2010.1126.
 1113
 1114 Radder, R. S. (2007) Maternally derived egg yolk steroid hormones and sex determination:
 1115 review of a paradox in reptiles, *Journal of Biosciences*, 32(6), pp. 1213–1220. doi:
 1116 10.1007/s12038-007-0123-z.
 1117
 1118 Reed, D. H. and Frankham, R. (2003) Correlation between Fitness and Genetic Diversity,
 1119 *Conservation Biology*, 17(1), pp. 230–237.
 1120
 1121 Refsnider, J. M., Bodensteiner, B. L., Reneker, J. L. and Janzen, F. J. (2013) Nest depth may not
 1122 compensate for sex ratio skews caused by climate change in turtles, *Animal Conservation*, 16(5),
 1123 pp. 481–490. doi: 10.1111/acv.12034.
 1124
 1125 Refsnider, Jeanine M., Warner, D. A. and Janzen, F. J. (2013) Does shade cover availability limit
 1126 nest-site choice in two populations of a turtle with temperature-dependent sex determination?,
 1127 *Journal of Thermal Biology*, 38(3), pp. 152–158. doi: 10.1016/j.jtherbio.2013.01.003.
 1128
 1129 Reneker, J. L. and Kamel, S. J. (2016) The maternal legacy: Female identity predicts offspring
 1130 sex ratio in the loggerhead sea turtle, *Scientific Reports*, 6: 29237. doi: 10.1038/srep29237.
 1131
 1132 Reusch, T. B. H. (2014) Climate change in the oceans: evolutionary versus phenotypically plastic
 1133 responses of marine animals and plants, *Evolutionary Applications*, 7, pp. 104–122. doi:
 1134 10.1111/eva.12109.
 1135
 1136 Rey, O., Eizaguirre, C., Angers, B., Baltazar-Soares, M., Sagonas, K., Prunier, J. G. and
 1137 Blanchet, S. (2020) Linking epigenetics and biological conservation: Towards a conservation
 1138 epigenetics perspective, *Functional Ecology*, 34(2), pp. 414–427. doi: 10.1111/1365-2435.13429.
 1139
 1140 Rhen, T. and Lang, J. W. (1995) Phenotypic plasticity for growth in the common snapping turtle:
 1141 Effects of incubation temperature, clutch , and their interaction, *The American Naturalist*, 146(5),
 1142 pp. 726–747.
 1143
 1144 Rhen, T. and Lang, J. W. (2004) Phenotypic effects of incubation temperature in reptiles, in
 1145 *Temperature-Dependent Sex Determination in Vertebrates*, pp. 90–98. doi: 10.1007/978-3-0348-

1146 6273-8_2.
 1147
 1148 Rhen, T. and Schroeder, A. L. (2017) The genetics of thermosensitive sex determination,
 1149 *Temperature*, 4(2), pp. 109–111. doi: 10.1080/23328940.2016.1259723.
 1150
 1151 Rhen, T. and Schroeder, A. L. (2010) Molecular mechanisms of sex determination in reptiles,
 1152 *Sexual Development*, 10(4), pp. 16-28. doi: 10.1159/000282495.
 1153
 1154 Rice, E. S., Kohno, S., St John, J., Pham, S., Howard, J., Lareau, L. F., O'Connell, B. L., Hickey,
 1155 G., Armstrong, J., Deran, A., Fiddes, I., Platt, R. N., Gresham, C., McCarthy, F., Kern, C., Haan,
 1156 D., Phan, T., Schmidt, C., Sanford, J. R., Ray, D. A., Paten, B., Guillette, L. J. and Green, R. E.
 1157 (2017) Improved genome assembly of American alligator genome reveals conserved architecture
 1158 of estrogen signaling, *Genome Research*, 27(5), pp. 686–696. doi: 10.1101/gr.213595.116.
 1159
 1160 Rimblot, F., Fretey, J., Mrosovsky, N. and Lescure, J. (1985) Sexual differentiation as a function
 1161 on the incubation temperature of eggs in the sea turtle *Dermochelys coriacea* (Vandelli, 1761),
 1162 *Amphibia Reptilia*, (83–92).
 1163
 1164 Rollinson, N. (2019) A simple explanation for the evolution and maintenance of temperature-
 1165 dependent sex determination, *Authorea*, doi: 10.22541/au.157592480.00795832.
 1166
 1167 Sanford, E. and Kelly, M. W. (2011) Local adaptation in marine invertebrates, *Annual Review of*
 1168 *Marine Science*, 3, pp. 509–535. doi: 10.1146/annurev-marine-120709-142756.
 1169
 1170 Santidrián Tomillo, P., Fonseca, L., Paladino, F. V., Spotila, J. R. and Oro, D. (2017) Are thermal
 1171 barriers “higher” in deep sea turtle nests?, *PLoS ONE*, 12(5), pp. 1–14. doi:
 1172 10.1371/journal.pone.0177256.
 1173
 1174 Santidrián Tomillo, P., Oro, D., Paladino, F. V., Piedra, R., Sieg, A. E. and Spotila, J. R. (2014)
 1175 High beach temperatures increased female-biased primary sex ratios but reduced output of
 1176 female hatchlings in the leatherback turtle, *Biological Conservation*, 176, pp. 71–79. doi:
 1177 10.1016/j.biocon.2014.05.011.
 1178
 1179 Santidrián Tomillo, P. and Spotila, J. R. (2020) Temperature-Dependent Sex Determination in
 1180 Sea Turtles in the Context of Climate Change: Uncovering the Adaptive Significance, *BioEssays*,

1181 2000146, pp. 1–6. doi: 10.1002/bies.202000146.

1182

1183 Sarre, S. D., Georges, A. and Quinn, A. (2004) The ends of a continuum: Genetic and

1184 temperature-dependent sex determination in reptiles, *BioEssays*, pp. 639–645. doi:

1185 10.1002/bies.20050.

1186

1187 Schofield, G., Hobson, V. J., Fossette, S., Lilley, M. K. S., Katselidis, K. A. and Hays, G. C.

1188 (2010) Fidelity to foraging sites, consistency of migration routes and habitat modulation of home

1189 range by sea turtles, *Diversity and Distributions*, 16, pp. 840–853. doi: 10.1111/j.1472-

1190 4642.2010.00694.x.

1191

1192 Schroeder, A. L., Metzger, K. J., Miller, A. and Rhen, T. (2016) A novel candidate gene for

1193 temperature-dependent sex determination in the common snapping turtle, *Genetics*, 203(May),

1194 pp. 557–571. doi: 10.1534/genetics.115.182840.

1195

1196 Schwabl, H. (1993) Yolk is a source of maternal testosterone for developing birds, *Proceedings of*

1197 *the National Academy of Sciences of the United States of America*, 90(December), p. 5. doi:

1198 10.1073/pnas.90.24.11446.

1199

1200 Sebastian, S. and Bulun, S. E. (2001) A Highly Complex Organization of the Regulatory Region

1201 of the Human CYP19 (Aromatase) Gene Revealed by the Human Genome Project. *The Journal*

1202 *of Clinical Endocrinology & Metabolism*, 86(10) 4600-4602

1203

1204 Senko, J., Mancini, A., Seminoff, J. A. and Koch, V. (2014) Bycatch and directed harvest drive

1205 high green turtle mortality at Baja California Sur , Mexico, *Biological Conservation*, 169, pp. 24–

1206 30. doi: 10.1016/j.biocon.2013.10.017.

1207

1208 Sheldon, B. C. and West, S. A. (2004) Maternal dominance, maternal condition, and offspring sex

1209 ratio in ungulate mammals, *American Naturalist*, 163(1), pp. 40–54. doi: 10.1086/381003.

1210

1211 Shen, Z. G. and Wang, H. P. (2014) Molecular players involved in temperature-dependent sex

1212 determination and sex differentiation in Teleost fish, *Genetics Selection Evolution*, 46(1), pp. 1–

1213 21. doi: 10.1186/1297-9686-46-26.

1214

1215 Shine, R. (1999) Why is sex determined by nest temperature in many reptiles?, *Trends in*

1216 *Ecology and Evolution*, 14(5), pp. 186–189. doi: 10.1016/S0169-5347(98)01575-4.

1217

1218 Shine, R., Elphick, M. J. and Donnellan, S. (2002) Co-occurrence of multiple, supposedly

1219 incompatible modes of sex determination in a lizard population, *Ecology Letters*, 5, pp. 486–489.

1220

1221 Simpson, E. R. (2004) Aromatase: Biologic Relevance of Tissue-Specific Expression. *Seminars*

1222 *in Reproductive Medicine*, 22(1), pp. 11-23.

1223

1224 Smith, C. A., Andrews, J. E. and Sinclair, A. H. (1997) Gonadal sex differentiation in chicken

1225 embryos: Expression of estrogen receptor and aromatase genes, *Journal of Steroid Biochemistry*

1226 *and Molecular Biology*, 60(5–6), pp. 295–302. doi: 10.1016/S0960-0760(96)00196-3.

1227

1228 Steele, A. L. and Warner, D. A. (2020) Sex-specific effects of developmental temperature on

1229 morphology, growth and survival of offspring in a lizard with temperature-dependent sex

1230 determination, *Biological Journal of the Linnean Society*, pp. 1–16. doi:

1231 10.1093/biolinnean/blaa038/5815733.

1232

1233 Stiebens, V. A., Merino, S. E., Roder, C., Chain, F. J. J., Lee, P. L. M. and Eizaguirre, C. (2013)

1234 'Living on the edge : how philopatry maintains adaptive potential', *Proceedings of the Royal*

1235 *Society*, 280, pp. 1–9.

1236

1237 Stocker, T. F., Qin, D., Plattner, G.-K., Alexander, L. V., Allen, S. K., Bindoff, N. L., Bréon, F.-M.,

1238 Church, J. A., Cubasch, U., Emori, S., Forster, P., Friedlingstein, P., Gillett, N., Gregory, J. M.,

1239 Hartmann, D. L., Jansen, E., Kirtman, B., Knutti, R., Kumar, K. K., Lemke, P., Marotzke, J.,

1240 Masson-Delmotte, V., Meehl, G. A., Mokhov, I. I., Piao, S., Ramaswamy, V., Randall, D., Rhein,

1241 M., Rojas, M., Sabine, C., Shindell, D., Talley, L. D., Vaughan, D. G. and Xie, S.-P. (2013)

1242 Technical Summary, in *Climate Change 2013: The Physical Science Basis. Contribution of*

1243 *Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*

1244 *Change*, pp. 33–109.

1245

1246 Strauss, J. F. and FitzGerald, G. A. (2018) *Steroid Hormones and Other Lipid Molecules Involved*

1247 *in Human Reproduction. Yen & Jaffe's Reproductive Endocrinology: Physiology,*

1248 *Pathophysiology, and Clinical Management: Eighth Edition.* Elsevier Inc. doi: 10.1016/B978-0-

1249 323-47912-7.00004-4.

1250

1251 Tanner, C., Marco, A., Martins, S., Abella-Perez, E. and Hawkes, L. (2019) Highly feminised sex
 1252 ratio estimations for the world's third-largest nesting aggregation of the loggerhead sea turtle,
 1253 *Marine Ecology Progress Series*, 621, pp. 209–219. doi: 10.3354/meps12963.

1254

1255 Telemeco, R. S., Abbott, K. C. and Janzen, F. J. (2013) Modeling the effects of climate change-
 1256 induced shifts in reproductive phenology on temperature-dependent traits., *The American*
 1257 *Naturalist*, 181(5), pp. 637–48. doi: 10.1086/670051.

1258

1259 Telemeco, R. S., Elphick, M. J. and Shine, R. (2009) Nesting lizards (*Bassiana duperreyi*)
 1260 compensate partly, but not completely, for climate change, *Ecology*, 90(1), pp. 17–22. doi:
 1261 10.1890/10-1922.1.

1262

1263 Tezak, B., Sifuentes-Romero, I., Milton, S. and Wyneken, J. (2020) Identifying sex of neonate
 1264 turtles with temperature-dependent sex determination via small blood samples, *Scientific*
 1265 *Reports*, 10, pp. 1–8. doi: 10.1038/s41598-020-61984-2.

1266

1267 Tilley, D., Ball, S., Ellick, J., Godley, B. J., Weber, N., Weber, S. B. and Broderick, A. C. (2019)
 1268 No evidence of fine scale thermal adaptation in green turtles, *Journal of Experimental Marine*
 1269 *Biology and Ecology*, 514–515, pp. 110–117. doi: 10.1016/j.jembe.2019.04.001.

1270

1271 Tomillo, P. S., Saba, V. S., Piedra, R., Paladino, F. V and Spotila, J. R. (2008) Effects of illegal
 1272 harvest of eggs on the population decline of leatherback turtles in Las Baulas Marine National
 1273 Park , Costa Rica, *Conservation Biology*, 22(5), pp. 1216–1224. doi: 10.1111/j.1523-
 1274 1739.2008.00987.x.

1275

1276 Trivers, R. L. and Willard, D. E. (1973) Natural selection of parental ability to vary the sex ratio of
 1277 offspring, *Science*, 179(4068), pp. 90–92.

1278

1279 Tuttle, J. and Rostal, D. (2010) Effects of nest relocation on nest temperature and embryonic
 1280 development of loggerhead sea turtles (*Caretta caretta*), *Chelonian Conservation and Biology*,
 1281 9(1), pp. 1–7. doi: 10.2744/CCB-0769.1.

1282

1283 Uhlenhaut, N. H., Jakob, S., Anlag, K., Eisenberger, T., Sekido, R., Kress, J., Treier, A. C.,
 1284 Klugmann, C., Klasen, C., Holter, N. I., Riethmacher, D., Schütz, G., Cooney, A. J., Lovell-Badge,
 1285 R. and Treier, M. (2009) Somatic sex reprogramming of adult ovaries to testes by FOXL2

1286 ablation, *Cell*, 139(6), pp. 1130–1142. doi: 10.1016/j.cell.2009.11.021.

1287

1288 Varela, M. R., Patrício, A. R., Anderson, K., Broderick, A. C., DeBell, L., Hawkes, L. A., Tilley, D.,

1289 Snape, R. T. E., Westoby, M. J. and Godley, B. J. (2019) Assessing climate change associated

1290 sea-level rise impacts on sea turtle nesting beaches using drones, photogrammetry and a novel

1291 GPS system, *Global Change Biology*, 25(2), pp. 753–762. doi: 10.1111/gcb.14526.

1292

1293 Visser, M. E. (2008) Keeping up with a warming world; assessing the rate of adaptation to climate

1294 change, *Proceedings of the Royal Society B: Biological Sciences*, 275(1635), pp. 649–659. doi:

1295 10.1098/rspb.2007.0997.

1296

1297 Wallace, B. P., Kot, C. Y., Dimatteo, A. D., Lee, T., Crowder, L. B. and Lewison, R. L. (2013)

1298 Impacts of fisheries bycatch on marine turtle populations worldwide: Toward conservation and

1299 research priorities, *Ecosphere*, 4(3), pp. 1–49. doi: 10.1890/ES12-00388.1.

1300

1301 Wang, D. S., Kobayashi, T., Zhou, L. Y., Paul-Prasanth, B., Ijiri, S., Sakai, F., Okubo, K.,

1302 Morohashi, K. I. and Nagahama, Y. (2007) Foxl2 up-regulates aromatase gene transcription in a

1303 female-specific manner by binding to the promoter as well as interacting with Ad4 binding

1304 protein/steroidogenic factor, *Molecular Endocrinology*, 21(3), pp. 712–725. doi: 10.1210/me.2006-

1305 0248.

1306

1307 Warner, D. A., Addis, E., Du, W. guo, Wibbels, T. and Janzen, F. J. (2014) Exogenous

1308 application of estradiol to eggs unexpectedly induces male development in two turtle species with

1309 temperature-dependent sex determination, *General and Comparative Endocrinology*, 206, pp.

1310 16–23. doi: 10.1016/j.ygcen.2014.06.008.

1311

1312 Warner, D. A., Mitchell, T. S., Bodensteiner, B. L. and Janzen, F. J. (2017) The effect of hormone

1313 manipulations on sex ratios varies with environmental conditions in a turtle with temperature-

1314 dependent sex determination, *Journal of Experimental Zoology Part A: Ecological and Integrative*

1315 *Physiology*, 327(4), pp. 172–181. doi: 10.1002/jez.2085.

1316

1317 Warner, D. A. and Shine, R. (2008) The adaptive significance of temperature-dependent sex

1318 determination in a reptile, *Nature*, 451(January), pp. 2006–2009. doi: 10.1038/nature06519.

1319

1320 Weber, S. B., Broderick, A. C., Groothuis, T. G. G., Ellick, J., Godley, B. J. and Blount, J. D.

1321 (2012) Fine-scale thermal adaptation in a green turtle nesting population., *Proceedings of The*
 1322 *Royal Society B*, 279(1731), pp. 1077–1084. doi: 10.1098/rspb.2011.1238.
 1323
 1324 Weishampel, J. F., Bagley, D. a. and Ehrhart, L. M. (2004) Earlier nesting by loggerhead sea
 1325 turtles following sea surface warming, *Global Change Biology*, 10(8), pp. 1424–1427. doi:
 1326 10.1111/j.1365-2486.2004.00817.x.
 1327
 1328 Wibbels, T., Bull, J. J. and Crews, D. (1991) Synergism between temperature and estradiol: a
 1329 common pathway in turtle sex determination?, *The Journal of experimental zoology*, 260(1), pp.
 1330 130–134. doi: 10.1002/jez.1402600117.
 1331
 1332 Wibbels, T. and Crews, D. (1994) Putative aromatase inhibitor induces male sex determination in
 1333 a female unisexual lizard and in a turtle with temperature-dependent sex determination, *Journal*
 1334 *of Endocrinology*, 141, pp. 295–299.
 1335
 1336 Wibbels, T., Rostal, D. and Byles, R. (1998) High pivotal temperature in the sex determination of
 1337 the olive ridley sea turtle, *Lepidochelys olivacea*, from Playa Nancite, Costa Rica, *Copeia*,
 1338 1998(4), pp. 1086–1088.
 1339
 1340 Witt, M. J., Hawkes, L. A., Godfrey, M. H., Godley, B. J. and Broderick, A. C. (2010) Predicting
 1341 the impacts of climate change on a globally distributed species: the case of the loggerhead turtle,
 1342 *Journal of Experimental Biology*, 213(6), pp. 901–911. doi: 10.1242/jeb.038133.
 1343
 1344 Wyneken, J., Epperly, S. P., Crowder, L. B., Vaughan, J. and Blair Esper, K. (2007) Determining
 1345 sex in posthatchling loggerhead sea turtles using multiple gonadal and accessory duct
 1346 characteristics, *Herpetologica*, 63(1), pp. 19–30. doi: 10.1655/0018-
 1347 0831(2007)63[19:dsipls]2.0.co;2.
 1348
 1349 Xia, Z.-R., Li, P.-P., Gu, H.-X., Fong, J. J. and Zhao, E.-M. (2011) Evaluating noninvasive
 1350 methods of sex identification in green sea turtle (*Chelonia mydas*) hatchlings, *Chelonian*
 1351 *Conservation and Biology*, 10(1), pp. 117–123. doi: 10.2744/CCB-0852.1.
 1352
 1353 Yntema, C. L. and Mrosovsky, N. (1982) Critical periods and pivotal temperatures for sexual
 1354 differentiation in loggerhead sea turtles, *Canadian Journal of Zoology*, 60(5), pp. 1012–1016. doi:
 1355 10.1139/z82-141.

1356

1357 Zbinden, J., Davey, C., Margaritoulis, D. and Arlettaz, R. (2007) Large spatial variation and
1358 female bias in the estimated sex ratio of loggerhead sea turtle hatchlings of a Mediterranean
1359 rookery, *Endangered Species Research*, 3, pp. 305–312. doi: 10.3354/esr00058.

1360

1361

1362 **Box 1: An overview of Aromatase**

1363 Aromatase is an enzyme encoded by the *Cyp19a1* gene (Strauss *et al.*, 2018), which is part of
1364 the cytochrome P450 superfamily. This protein is the only known enzyme to catalyse the
1365 conversion of androgens to estrogens, a process that occurs throughout all vertebrate taxa. The
1366 P450 superfamily is an ancient lineage of genes that diverged early in the evolution of vertebrates
1367 (Simpson, 2004; Boon *et al.*, 2012; Nelson *et al.*, 2013), with gonadal synthesis of estrogens
1368 originating 500 mya (Lange *et al.*, 2002). *Cyp19a1* is highly conserved (Conley *et al.*, 2001). In
1369 fish and reptiles, there are two aromatase isomorphs encoded by the *Cyp19a1* and *Cyp19b1*
1370 genes and expressed in the gonads and brain respectively (Boon *et al.*, 2012). For most species,
1371 the majority of estrogen biosynthesis occurs in the gonads, with biosynthesis in the brain
1372 associated with behaviour (Simpson, 2004). In comparison, a single gene encodes human
1373 aromatase, with tissue specific promoter regions found on exon 1 of this gene, enabling its
1374 biosynthesis to occur in a greater range of tissues (Sebastian *et al.*, 2001; Bulun *et al.*, 2004).
1375 The up-regulation of *Cyp19a1* in the gonads is required for ovarian differentiation in fish (Guiguen
1376 *et al.*, 2010), birds (Smith *et al.*, 1997) and reptiles (Jeyasuria *et al.*, 1998), but not in mammals,
1377 where the knockout of *Cyp19a1* does not prevent ovaries from developing (Fisher *et al.*, 1998).

1378

1379

1380

1381

Accepted Article

Species	Location	T _{piv} (°C)	Source of T _{piv}
Green (Esteban <i>et al.</i> , 2016)	Chagos Archipelago	29	Review (Ackerman, 1997)
Green (Broderick <i>et al.</i> , 2001)	Ascension Island	29	Review (Ackerman, 1997)
Green (Booth <i>et al.</i> , 2006)	Heron Island, Australia	27.5	T _{piv} previously calculated for population (Booth <i>et al.</i> , 2001)
Hawksbill (Esteban <i>et al.</i> , 2016)	Chagos Archipelago	29	Review (Ackerman, 1997)
Hawksbill (Glen <i>et al.</i> , 2004)	Antigua	29.2	Review (Mrosovsky <i>et al.</i> , 1991)
Leatherback (Santidrián Tomillo <i>et al.</i> , 2014)	Costa Rica	29.4	T _{piv} previously calculated for same population (Binckley <i>et al.</i> , 1998)
Loggerhead (Laloë <i>et al.</i> , 2014)	Cabe Verde	29, 28.8 and 29.2	Mathematical modelling - Fit three different T _{piv} and kept 29 °C
Loggerhead (Zbinden <i>et al.</i> 2007)	Zakynthos	29.3	T _{piv} previously calculated for same population (Mrosovsky <i>et al.</i> , 2002)
Loggerhead (Öz <i>et al.</i> , 2004)	Turkey	29	T _{piv} previously calculated for same population (Kaska <i>et al.</i> , 1998)
Loggerhead (Hanson <i>et al.</i> , 1998)	Florida	29	T _{piv} previously calculated for same population (Mrosovsky, 1988)
Loggerhead (Tanner <i>et al.</i> , 2019)	Cabo Verde	29.25	T _{piv} previously calculated for different population (Marcovaldi <i>et al.</i> , 1997)
Loggerhead (Jribi <i>et al.</i> , 2014)	Tunisia	29.7	T _{piv} previously calculated for different population (Mrosovsky <i>et al.</i> , 2002)





